
Safety Assessment of Phytosterols as Used in Cosmetics

Status: Final Report
Release Date: January 27, 2014
Panel Meeting Date: December 9-10, 2013

The 2013 Cosmetic Ingredient Review Expert Panel members are: Chairman, Wilma F. Bergfeld, M.D., F.A.C.P.; Donald V. Belsito, M.D.; Curtis D. Klaassen, Ph.D.; Daniel C. Liebler, Ph.D.; Ronald A. Hill, Ph.D.; James G. Marks, Jr., M.D.; Ronald C. Shank, Ph.D.; Thomas J. Slaga, Ph.D.; and Paul W. Snyder, D.V.M., Ph.D. The CIR Director is Lillian J. Gill, D.P.A. This report was prepared by Lillian C. Becker, Scientific Analyst/Writer.

© Cosmetic Ingredient Review

1101 17th Street, NW, Suite 412 ∇ Washington, DC 20036-4702 ∇ ph 202.331.0651 ∇ fax 202.331.0088 ∇ cirinfo@cir-safety.org

ABSTRACT

Phytosterols functions in cosmetics include skin-conditioning agents, hair conditioning agents, viscosity increasing agents, skin protectants, antioxidants, and fragrances. The Panel reviewed relevant animal and human data related to these ingredients including results of tests for relevant estrogenic activity. Industry should use good manufacturing practices to limit impurities. The Panel concluded that phytosterols were safe in the present practices of use and concentration described in this safety assessment.

INTRODUCTION

This report reviews the available scientific information relevant to the safety of a group of 26 phytosterols and steryl alkanoates as used in cosmetics. The functions of these ingredients include: skin-conditioning agents, hair conditioning agents, viscosity increasing agents, skin protectants, antioxidants, and fragrances (Table 1).¹ The ingredients in this report are:

- brassica campestris (rapeseed) sterols
- canola sterols
- C10-40 isoalkyl acid phytosterol esters
- dihydrophytosteryl octyldecanoate
- euterpe oleracea sterols
- glycine soja (soybean) sterols
- perseae gratissima (avocado) sterols
- phytosterols
- phytosteryl butyrate
- phytosteryl canolate
- phytosteryl caprylate/caprinate
- phytosteryl hydroxystearate
- phytosteryl isostearate
- phytosteryl linoleate
- phytosteryl linoleate/linolenate
- phytosteryl macadamiate
- phytosteryl nonanoate
- phytosteryl oleate
- phytosteryl rice branate
- phytosteryl ricinoleate
- phytosteryl sunflowerseedate
- punica granatum sterols
- beta-sitosterol
- beta-sitosteryl acetate
- soy sterol acetate
- tall oil sterol

Plant sterols, or phytosterols, occur naturally as free alcohols and as fatty acid esters (i.e., naturally occurring steryl alkanoates). The ingredients in this report are sterol alcohols or esters (in some cases mixtures of both) extracted from plants, which in some cases have been saponified to the free alcohols and then esterified with plant-derived fatty acids. The resultant ester-derivatized phytosterols (i.e., steryl alkanoates) share substantial component overlap with the naturally-occurring phytosterol esters. Most of these derived esters are synthetic analogs of the components of the naturally occurring phytosterol esters. Because there is a large amount of component overlap among the ingredients in this group, these ingredients can be reviewed as an ingredient family. The structural similarities among the compounds or components (i.e., phytosterols and fatty acid), the similarities of their physicochemical properties, and functions and concentrations of these ingredients in cosmetics enable grouping these ingredients and reading across the available toxicological data to support the safety assessment of the entire group. Table 2 lists the component compounds, noting whether or not these components are cosmetic ingredients and whether or not they have been reviewed by the Cosmetic Ingredient Review (CIR) Expert Panel (Panel). All of the reviewed component cosmetic ingredients were found to be safe or safe with qualifications. Butyric acid, caprylic acid/capric acid, and linoleic acid/linolenic acid have not been reviewed. Octyldecanoic acid is not a cosmetic ingredient.

In 2000, the Panel found the data on PEG-5, -10, -16, -25, -30, and -40 soy sterols to be insufficient to support the safety of these ingredients.² In 2004, the Panel found these PEG soy sterols to be safe as used in an amended safety assessment that included data on phytosterols and phytosterol esters.³ The Panel's approach in these safety assessments was to review the safety of PEGs and phytosterols/soy sterols, as well as the conjugated polyethers, and to assess the safety of the PEG phytosterols from those data. Because the data on the phytosterols/soy sterols are relevant for this safety assessment, summaries of the phytosterols/soy sterols data from these two safety assessments are provided in the appropriate sections.

Many of the phytosterols in this study are from edible plant sources. Exposure to these phytosterols from consuming foods results in much greater systemic doses than could result from the use of cosmetic products. It was noted in the PEG soy sterol reports that phytosterols and phytosterol esters are not significantly absorbed after oral exposure, and thus, did not result in systemic exposure.^{2,3} Therefore, acute and repeated dose oral toxicity potential of these phytosterols as cosmetic ingredients will not be addressed again in depth in this report. The focus of this report is on: reproductive toxicity, genotoxicity, carcinogenicity, dermal irritation, and sensitization.

CHEMISTRY

Definition, Structure, and Composition

See Table 1 for information on definitions and functions.

The phytosterol ingredient group is comprised of the plant-derived free sterols and their esters, the steryl alkanooates. β -Sitosterol is an example of a discrete, free phytosterol ingredient.

To generate a steryl alkanooate with an ester at the 3-position of the sterol, the hydroxyl group at the 3-position of the cyclopentenophenanthrene scaffold is esterified by reacting with an alkyl acid or acid chloride (i.e., β -sitosteryl acetate). Representative phytosterols are also provided in Figure 1.




Figure 1. Examples of discrete molecules found in phytosterol ingredients

Phytosterols occur in plants in the free alcohol, steryl alkanooate, or glycoside forms. The free phytosterols are characteristic components of the non-saponifiable fractions of plant oils.⁴ The steryl alkanooate and glycoside forms, however, are broken down to the free phytosterol form (and respective acid or sugar) under saponification conditions. The majority of the ingredients in this report are mixtures of either sterols or steryl alkanooates, with component concentrations that vary with growth and extraction conditions.

As one example, soybean oil that had been alkali-refined typically contained 0.446 mg/100 mg oil of total sterols

and 0.287 mg/100 mg oil of free sterol. The ratio of esterified to free sterol was 0.55.⁵

Palm oil/crude palm oil contains 300 – 620 ppm phytosterols, 60% of which is beta-sitosterol with 38% being stigmasterol and campesterol.⁶

Product information on refined soy sterols reported that they contain ~ 88% total sterol content. Of that percentage, 56% is γ -sitosterol, 28% is campesterol, and 4% is stigmasterol. Other compounds isolated from the phytosterols are 4% - 6% sterol hydrocarbons and cholesterol, and 4% - 6% triterpene alcohols, keto-steroids, and other steroid-like substances.⁷

A typical palm phytosterol is characterized as a crystalline waxy powder or a free-flowing granular powder that is white to off white and practically odorless.^{6,8} Its melting point is reported to be 131-141°C. The phytosterol content is at least 80% (50% beta sitosterol; 20% campesterol; and 2% stigmasterol). Palm-derived phytosterols were reported to be stable for 36 months when stored in a cool dry environment in unopened containers. The product is sensitive to air, light, and heat.

PEGs Soy Sterol Report

The chemical characterization of a plant sterol material is provided in Table 3. The distributions of phytosterols in common vegetable oils are provided in Table 4.

In an analysis of another source of phytosterols (source not provided), it was reported that the principal phytosterols were present as follows: β -sitosterol, 47.9%; campesterol, 28.8%; and stigmasterol, 23.3%. In an analysis of phytosterol esters, it was reported that the principal phytosterols were present as fatty acid esters: β -sitosterol, 47.3%; campesterol, 28.1%; and stigmasterol, 24.5%. The distribution of the fatty acid chain lengths was consistent with fatty acids derived from sunflower oil.^{9,10}

Physical and Chemical Properties

Physical and chemical properties of representative phytosterols in the form of vegetable oil sterols and tall oil sterols are provided in Table 5.

Phytosterols and their fatty acid esters are thermally stable and degrade only at high temperatures (>100°C) in the presence of oxygen.¹¹

Method of Manufacture

Free phytosterol alcohols and phytosterol alkanolates are characteristic components of plant oils; saponification of these oils is the primary means of producing free phytosterol alcohols for commercial use.⁴

Soy sterol is isolated from soybean oil distillates in a saponification process in which the phytosterol alcohols are separated from the fatty acids by extraction with an organic solvent.² The phytosterols in the resulting extract are separated from the tocopherols in the mother liquor, and then purified and/or separated into constituent sterols.

Tall oil sterol, an example of a phytosterol mixture, is obtained from tall oil soap in a multi-step process.⁴ The production process involves fractional distillation of the tall oil soap to remove volatile compounds. The resulting residue (tall oil pitch), containing esterified sterols (i.e., steryl alkanolates), is treated with alkali (saponified) to release the free sterol alcohols. After neutralization, the material is subjected to a two-stage distillation process. The distillate is then dissolved in methanol/methylethylketone solvent and the sterols crystallizing from this solution are obtained by filtration, washed with solvent and dried. This procedure results in a lower stanol and a higher sterol content of the phytosterol mixture. Conifers that have naturally lower stanol content are now used as the primary source of the tall oil soap. Stanols (obtained by catalytic hydrogenation of the phytosterol mixture) are added before the crystallization step to maintain the original stanol/sterol ratio. The phytosterol composition of the tall oils produced from the two processes is provided in Table 6.

Steryl alkanolates are produced from free sterols by classical esterification methods, using acids or acid chlorides. Sterol alkanolates may be derived from neutralized, refined, bleached and deodorized (N/RBD) reaction with soybean distillates.^{12,13} Crude soybean oil is degummed, neutralized, bleached and deodorized to yield N/RBD soybean oil and distillates. The deodorized distillate undergoes further processing (crystallization and/or distillation), resulting in a sterol mixture. This sterol mixture is then crystallized and esterified with fatty acids (from food grade vegetable oils such as rapeseed or sunflower oil), washed, bleached and deodorized to give the final plant steryl alkanolates.

A manufacturer reported that for the manufacture of glycine soja (soybean) sterols, and punica granatum sterols, the raw materials are tested for acceptable qualifications (not specified) before they are cold pressed for oil.^{14,15} The oil is then tested for quality (not specified) before the oil is fractionated to isolate the sterols. Pomegranate sterols are heat-sterilized at 100°C before fractionation.¹⁶

A manufacturer reports that the extraction process for palm oil phytonutrients is an integrated process for the recovery of phytosterols as well as vitamin E and squalene.⁸ The process comprises the steps of acid/alkaline catalyzed esterification/transesterification process of palm oil with a lower alkyl alcohol, multi-stage vacuum distillation of alkyl esters, saponification of the phytonutrients concentrate, crystallization of phytosterols, and partitioning of vitamin E and squalene with organic solvents.

Impurities

Commercial tall oil sterols/stanols were reported to contain < 0.1 mg/kg lead.¹¹ Commercial vegetable oil sterols,

in general, were reported to have < 2.0 mg/kg impurities (mercury, < 0.1%; lead, < 0.1%; cadmium, < 0.1%; and arsenic, < 0.1%). Both were reported to contain < 2 ppb PAHs and < 1.5 ng-TEQ/kg dioxins and dioxin-like PCBs. No pesticides were detected.

In an analysis of *Euterpe oleracea* sterols, glycine soja (soybean) sterols, and *Punica granatum* sterols, none of these ingredients contained detectable levels of an array of potential allergens, including amyl cinnamal, benzyl alcohol, citronellol, coumarin, linalool, and farnesol (Table 7). Another analysis did not detect any of several pesticides, including dichlorodiphenyltrichloroethane (DDT; detection level 1.00 mg/kg), methidathion (0.20 mg/kg), and pyrethrins (3.00 mg/kg).²⁰⁻²²

Phytosterols derived from palm phytosterol are reported to typically meet the following parameters: mercury, < 1 ppm; cadmium, < 1 ppm; arsenic, < 1 ppm; lead, < 1 ppm; benzo(a)pyrene, < 2 ppm.⁸ They are allergen free with regards to peanut, eggs, milk/milk by-products, wheat or gluten, tree nuts, soy, fish and shellfish. Assays for pesticide impurities are reported as follows: araquat, <0.05 ppm; diquat, <0.05 ppm; DDT, <0.05 ppm, hexachlorocyclohexane, <0.05 ppm. These pesticides are known to be used in palm oil plantations in Malaysia.

In assessing the data on soybean oil sterols, the Scientific Panel on Dietetic Products, Nutrition and Allergies noted that there are limited analytical data of sufficient sensitivity and reliability regarding the possible residual allergen (protein) content of phytosterols.¹² Specifically, the limited analytical data regarding the protein (allergen) content of N/RBD soybean oil-derived plant stanol esters were insufficient to predict the likelihood of adverse reactions in soybean-allergic individuals. This panel concluded, however, that because the starting material is refined soybean oil and there is an adequate subsequent production process, it is not very likely that this product will retain enough residual protein to cause a severe allergic reaction in the majority of soybean-allergic individuals.

The final protein content of N/RBD soybean oils (the source of soy phytosterols) is known to depend on the quality and efficiency of purification steps.¹² The protein content of N/RBD oils may be reduced to low levels within the 0.02-0.44 µg/kg range.¹⁷ When two samples of edible soy oil (crude virgin and deodorized) were analyzed for proteins, 1.89 µg/mL and 0.32 µg/mL proteins were found in the samples.¹⁸

When selected phytosterol samples (a phytosterol blend and a phytosterol blend spiked with reference protein) were analyzed for residual soybean protein using enzyme-linked immunosorbent assay (ELISA), soy protein was not detectable at or above the 10-20 µg/g detection limit.¹⁹

All hydrophilic extracts of vegetable oil deodorized distillate (VOD) samples (n = 9) analyzed by immunoblotting with soy-specific antiserum from rabbits and by IgE-immunoblotting with a pooled human serum detected no soy protein or other protein.¹⁹ There was no IgE binding with the VOD or the phytosterol samples using either the pooled human serum or the serum of one subject who had experienced mild oral allergy syndrome after a double-blind placebo controlled food challenge (DBPCFC) with phytosterols. The authors concluded that no IgE-binding proteins were present in the VOD and phytosterol samples at or above 1 and 10 µg/g, respectively.

Refined soybean oils exhibited no detectable IgE binding activity using immunoblotting and enzyme allergosorbent test (EAST) inhibition assays.²³

PEGs Soy Sterol Report

Analyses of various lots of soy sterols for pesticide residues were negative for a number of pesticides, including polychlorinated biphenyls (PCB), dichlorodiphenyldichloroethylene (DDE), DDT, malathion, and β-hexachloride.²⁴ In an analysis of phytosterols (source not provided), no impurities were found.^{9,10}

USE Cosmetic

Data on ingredient usage are provided to the Food and Drug Administration (FDA) Voluntary Cosmetic Registration Program (VCRP; Table 8).²⁵ A survey was conducted by the Personal Care Products Council (Council) of the maximum use concentrations for ingredients in this group.²⁶

Data were available from both the VCRP and the Council for the following ingredients:

- Brassica campestris (rapeseed) sterols was reported to be used in 50 leave-on products and 7 rinse-off products. Leave-on products were reported to contain up to 7% (the highest amount in lipstick) brassica campestris (rapeseed) sterols and 0.13% in rinse-off products.
- Glycine soja (soybean) sterols was reported to be used in 194 leave-on products (mostly skin care and makeup products), 45 rinse-off products, and one bath product. Leave-on products were reported to contain up to 1% (the highest concentration in eye lotion, cuticle softeners, and other skin preparations) and up to 4.1% in rinse-off products (the highest concentration in skin cleansing products). This ingredient was reported to be used in tonics, dressings and other hair grooming aids, including an aerosol and a pump spray at 0.000001%.
- Phytosterols was reported to be used in 177 leave-on products. It was reported that phytosterols was used in lipsticks up to 5%, non-spray deodorants up to 0.06%, and eye makeup up to 2%. It is also used in 215 rinse-off products. It was reported that phytosterols was used in hair products up to 2.4%, bath soaps and detergents up to 0.005%, and indoor tanning preparations up to 0.0001%. It was also reported to be used in face powders up to 0.05%.

- Phytosteryl isostearate was reported to be used in 15 leave-on products and one rinse-off product. This ingredient was reported to be used up to 3% in leave-on products and in rinse-off products up to 0.5%. It is used in lipsticks up to 3% and in eye makeup up to 0.5%.
- Phytosteryl [phytosterol] macadamiate was reported to be used in 181 leave-on products (100 lipsticks) and in two rinse-off products. It was reported to be used in leave-on products up to 8% and in rinse-off products up to 1%. It was reported to be used in lipsticks up to 7% and in moisturizing products up to 8%.
- Phytosteryl oleate was reported to be used in 20 leave-on products (including 6 paste masks/mud packs). It was reported to be used up to 3%.
- Phytosteryl rice branate was reported to be used in an eye makeup and a moisturizing product. The Council reported that it was used in eye lotions up to 1%, foundations up to 0.5%, and face and neck products up to 0.5%.
- Punica granatum sterols was reported to be used in 29 rinse-off products and in two rinse-off products. It was reported to be used up to 5% in leave-on products (including lipsticks).
- Beta-sitosteryl was reported to be used in 46 leave-on products and in two rinse-off products. It was reported to be used in leave-on products up to 0.06%.
- Tall oil sterol was reported to be used in 7 leave-on products. It was reported to be used up to 0.0046%, including in skin cleansing products up to 0.0006%.

Data were available only on the frequencies of use (VCRP) for the following ingredients:

- Euterpe oleracea sterols was reported to be used in one lipstick and one foundation.
- Soy sterol acetate was reported to be used in one moisturizing product.

Data were available only on use concentrations (Council) for the following ingredients:

- Persea gratissima (avocado) sterols was reported to be used in eye lotion up to 1%, lipstick up to 0.65%, and face and neck products up to 0.1%.
- Phytosteryl canolate was reported to be used in eye shadow up to 0.06%.

There were no uses or concentration of use data reported for:

- canola sterols
- C10-40 isoalkyl acid phytosterol esters
- dihydrophytosteryl octyldecanoate
- phytosteryl butyrate
- phytosteryl caprylate/caprinate
- phytosteryl hydroxystearate
- phytosteryl linoleate
- phytosteryl linoleate/linolenate
- phytosteryl nonanoate
- phytosteryl ricinoleate
- phytosteryl sunflowerseedate
- punica granatum sterols
- beta-sitosteryl acetate

As noted above, uses were reported for glycine soja (soybean) sterols in propellant and pump spray tonics, dressings and other hair grooming aids up to 0.000001%, and phytosterols are reported to be used in face powders up to 0.05%. In practice, 95% to 99% of the droplets/particles released from cosmetic sprays have aerodynamic equivalent diameters >10 µm, with propellant sprays yielding a greater fraction of droplets/particles below 10 µm compared with pump sprays. Therefore, most droplets/particles incidentally inhaled from cosmetic sprays would be deposited in the nasopharyngeal and bronchial regions and would not be respired (i.e., they would not enter the lungs) to any appreciable extent.²⁷⁻³²

Non-Cosmetic

Phytosterols (stigmasterol-rich plant sterols: stigmasterol, >85%; brassicasterol, 1.7%; β-sitosterol, 3%; campesterol, 1.7%) are used in ready-to-freeze alcoholic beverages as a stabilizer.³³

Phytosterols and phytostanols are commonly used in food products for their property of absorption reduction of cholesterol in the gut and for lowering cholesterol content in food products.¹¹ The optimal daily intake for the former purpose is 2 - 3 g. For example, in Europe, phytosterol esters are added to margarines and low-fat spreads (3.4 g/30 g), yogurts (1.25 g/125 mL), yogurt drinks (3.4 g/100mL), and milk (5 g/L).

When making a claim of the heart health benefits of consuming plant sterol/stanol esters for food products, the FDA requires that there be at least 80% β-sitosterol, campesterol, and stigmasterol (combined weight). The daily dietary intake levels of plant sterol esters of 1.3 g/d and 3.4 g/d for plant stanol esters (spread out in at least two servings at different times of the day with other food) are associated with reduced risk of coronary heart disease.[101.CFR101.83]

Suggested uses for phytosterols from palm oil were: dietary supplements, functional ingredient for margarine/butter/oil, an emulsifier, and feed additives for prawns, lobsters, etc.⁶

The Scientific Committee on Food (SCF) and European Food Safety Authority (EFSA) concluded that phytosterols, phytostanols and their esters could safely be approved for use in various foods (i.e., yellow fat spreads, soya drinks, salad dressings, rye bread) within the EU at levels resulting in intake of up to 3 g/day.^{12,13,19,34-45}

TOXICOKINETICS

Absorption, Distribution, Metabolism, and Excretion

No published dermal or inhalation ADME studies were discovered and no unpublished data were submitted.

Oral

The Western diet consists of ~160-360 mg/d phytosterols consisting of ~80% β -sitosterol. The diet also includes some campesterol and stigmasterol, small amounts of brassicasterol, and trace amounts of Δ -5-saturated plant stanols.⁴⁶

Less than 5% of dietary phytosterols, phytostanols, and their esters are absorbed in the gastrointestinal tracts of rats and humans.³³ Following absorption, phytosterols/phytostanols are transported in the serum via high density lipoproteins (HDL) in rats and low density lipoproteins (LDL) in humans to various organs and tissues, mostly to the liver. In the liver, phytosterols may be converted to bile acids. Absorbed phytosterols and phytostanols are predominantly excreted as such or as bile acids by the biliary route into the feces. The metabolic fate of phytosterols, phytostanols, and their esters is similar between rats and humans. The individual plant sterols are metabolized in a similar manner to each other. The phytosterols that are not absorbed in the gastrointestinal tract, or excreted as such in the bile, continue to the colon intact and are excreted in the feces.⁴⁶⁻⁴⁹

In an oral study (n = 10 healthy men), the average intestinal absorption of individual phytosterols were: campesterol, 9.6%; stigmasterol, 4.8%; and sitosterol, 4.2%.⁵⁰ The authors noted that these results were consistent with the results of animal studies showing that, for analogs of cholesterol, increasing the side-chain length of cholesterol reduced the absorbability of the sterol, with the exception of campesterol. The 5 α -campesterol-saturated (the corresponding stanol) had greater absorbability than campesterol. Absorption was measured by an intestinal perfusion technique over a 50-cm segment of the upper jejunum.

In another study using male subjects, the biliary secretion rate of β -sitosterol was faster (1.23 mg/h) than that of campesterol (0.76 mg/h).⁵¹

A female subject excreted increasing amounts of β -sitosterol, campesterol and stigmasterol through the skin as oral intake of phytosterols increased over sustained periods of time.⁵² When phytosterols were removed from the diet, the amount of β -sitosterol in the skin decreased from 6 mg/d to 0.08 mg/d within 83 days and finally became undetectable. Similar results were reported for the other two phytosterols. Twenty days after the administration of 30 g/d phytosterols, β -sitosterol, as well as campesterol and stigmasterol, reappeared in the skin and was excreted at 5 mg/d by 6 weeks.

Plant sterols, including stigmasterol and stanols (34 g/kg in feed), were able to cross the blood-brain barrier in a 90-day feeding study of Watanabe heritable hyperlipidemic rabbits.⁵³

Cytotoxicity

β -sitosterol (200 μ g/mL in ethanol) and β -sitosterol/campesterol (50%/40%; 200 μ g/mL in ethanol) were cytotoxic to mouse macrophages (strain C57BL/6).⁵⁴ Cytotoxicity was demonstrated through cell viability, lipid uptake, lactate dehydrogenase (LDH) leakage, cellular protein content, and a 3'-[1-(phenylaminocarbonyl)-3,4-tetrazolium]bis(4-methoxy-6-nitro)benzene sulfonic acid hydrate (XTT) assay.

Phytosterols (0.01 – 40 mM; derived from pomegranate) were not cytotoxic to HepG2 cells in a Neutral Red Cytotoxicity assay.⁵⁵

PEGs SOY STEROL REPORT

β -Sitosterol (100 μ g/ml; 5% in dimethyl sulfoxide (DMSO) and saline) was cytotoxic to seven cancer cell lines.⁵⁶

Sitosterolemia

Sitosterolemia (also referred to as phytosterolemia) is a rare autosomal recessively inherited lipid metabolic disorder that is characterized by hyperabsorption and decreased biliary excretion of dietary sterols. This condition leads to hypercholesterolemia, tendon and tuberous xanthomas, premature development of atherosclerosis, and abnormal hematologic and liver function test results.^{57,58}

ANIMAL TOXICOLOGY

Many of the phytosterols in this report are from edible sources; exposure to these phytosterols from food would presumably result in much larger systemic exposure than those resulting from use in cosmetic products. A summary of toxicity data on phytosterols, including oral data, from the PEG soy sterol report is presented below. However, this report does not focus on oral toxicity, but instead on the potential for reproductive toxicity, genotoxicity, carcinogenicity, irritation and sensitization via routes of exposure consistent with currently reported uses in cosmetics. A summary of toxicity data on phytosterols, including oral data, from the PEGylated soy sterol report is presented below for background information on toxicity.

Dermal - Non-Human

The dermal LD₅₀ of two mixtures of phytosterol esters was reported to be > 2000 mg/kg.⁵⁹ A wood-derived mixture (a stanol composition ~94% β -sitostanol and ~6% campestanol in corn oil; WDPSE) and a vegetable oil-derived mixture of

phytostanol esters (~68% β -sitosterol and ~32% campestanol in corn oil; VODPSE) were administered dermally to rats (n = 5/sex) for 24 h in accordance to the Organization for Economic Co-operation and Development (OECD) Test Guideline 404. No deaths or clinical signs of toxicity were observed after application of WDPSE. One male rat in the VODPSE group died of unrelated causes during the 14-day observation period.

PEGs Soy Sterol Report

Wistar rats administered a basal diet supplemented with cholesterol and maize phytosterols (72.5% β -sitosterol, 0.5% campesterol, and 7% stigmasterol) had decreased hepatic cholesterol concentrations.⁶⁰ Rats administered the high dose of cholesterol and phytosterols had decreased malic enzyme and acetylCoA carboxylase activities, and had hypotriglyceridemia.

Wistar rats administered subcutaneous injections of 2.5 to 5 μ g/1 g β -sitosterol for 60 days had no gross or microscopic lesions of the liver or kidneys.⁶¹ Rats administered 10 μ g/1 g had mild fibroblastic proliferation around the hepatic lobules and mild microscopic lesions of the kidney. Serum cholesterol was reduced in a dose-dependent manner, and serum protein was markedly reduced in rats of the high dose group.

In a 90-day oral toxicity study in female Wistar rats (n = 4), diets containing plant phytosterol esters up to 8.1 % were well tolerated.⁶² Some small hematology and blood chemistry variations from the controls were observed. No treatment related effects were observed with organ weights and histological examination and there was no evidence of systemic toxicity. Absent any organ effects, the small hematology and blood chemistry variations were not considered of toxicological significance.

Thirteen dogs fed a basic diet supplemented with 0.5 to 1.0 g/kg/day β -sitosterol had no gross or microscopic changes after 8 to 22 months of treatment. Weight gains and clinical parameters did not differ from controls.²⁴

No adverse effects or gross or microscopic abnormalities were observed in six New Zealand white rabbits of both sexes that were given feed containing 3% cottonseed sterols and 4% soy sterols for 70-212 days.²⁴

REPRODUCTIVE AND DEVELOPMENTAL TOXICITY

In a two-generation feeding study, the no observed adverse effect level (NOAEL) for phytosterol esters was \geq 8.1% (the highest dose tested) in the diet.^{63,64} Wistar rats, F₀ generation, (n = 28/sex) were administered phytosterol esters (0, 1.6%, 3.2%, 8.1%) in feed for 10 weeks before mating, and continuing through gestation and weaning. The F₁ generation (n = 28/sex) were fed the same diet as their F₀ parents and mated after 10 weeks. The analysis of the phytosterols revealed the following breakdown: brassicasterol (2.9%), campesterol (26.7%), stigmasterol (17.7%), β -sitosterol (51.0%), cholesterol (0.2%), and unknowns (1.5%).

There were no maternal or teratogenic effects attributed to the test substance. There were no effects on fertility and reproductive parameters, including sexual maturity, estrous cycle length, precoital time, and the histopathology of reproductive tissues in either generation. There were no developmental or reproductive effects observed in either generation. Necropsies were unremarkable.

In the F₁ generation, the body weights of male rats in the low-dose group were decreased in weeks 6 - 9 and 11 - 14. In the high-dose groups of the F₀ and F₁ generations, body weights of the male rats were reduced when compared to the body weights of control group males. Decreased weights were observed in weeks 2 - 7 and on occasions during weeks 5 - 15 and 18 - 19. No differences in the times to reach preputial separation were observed among F₁ males. In F₀ males, statistically significant increase in the relative liver weights (low- and mid-dose groups), increase in the relative adrenal weights (low dose group), and statistically significant increase in relative brain weights (low- and high-dose groups) in F₁ males were considered small changes and not dose-related so were considered to be of no biological or toxicological significance.

The NOAEL (8.1%) is equivalent to 3.3 - 6.5 g phytosterol esters/kg/d during the 10-week pre-mating period (~ 2.1 - 4.1 g phytosterols/kg/d or 400 - 900 mg stigmasterol/kg/d) and 2.5 - 9.1 g phytosterol esters/kg/d during gestation (~1.4 - 5.7 g/kg/d or 300 - 1200 mg stigmasterol/kg/d). The authors concluded that 2.5 - 9.1 g phytosterol esters/kg/d and 1.54 - 5.62 g phytosterols/kg/d (~ 335 - 1219 mg stigmasterol/kg/d), dependent on the phase of the study, was the NOAEL of daily oral administration of phytosterol esters for two successive generations.^{63,64}

There were no signs of reproductive toxicity to American minks (n = 70/sex) orally administered β -sitosterol (at 0, 5, 10 or 50 mg/kg/d) for 10 months.⁶⁵ In the second part of the study, after 7 months of exposure, males (n = 10, 11) were mated with 4 - 5 females each. There were no differences in number of pregnant females, litter and kit numbers, postnatal mortality and development and there were no treatment-related changes. After 3 months of exposure, 15 males/group were killed and investigated for organ weights and hematological and clinical chemistry parameters. Males exhibiting low quality fur were selected for this part of the study. There were differences in body fat masses (omental, mesenteric, retroperitoneal, intra-abdominal fat) reported, but increases in fat masses were not dose dependent. There were increased blood hemoglobin and serum high-density lipoprotein cholesterol concentrations observed.

Subcutaneous injections of β -sitosterol (5 mg/kg/d) for 16 to 48 days decreased sperm concentrations and fertility, and decreased testis and accessory sex tissue weights in a time-dependent manner in male Wistar rats.⁶⁶ Rats administered 0.5 mg/kg/d had a decrease in sperm concentration of the caput epididymis after 48 days of treatment, but no reduction in fertility. The observed decreases in sperm concentration persisted after termination of treatment, and appeared to be due to a reduction in the rate of spermatogenesis.

TESTS FOR ESTROGENIC ACTIVITY

In Vitro

There were no signs of estrogenic activity of phytosterols and phytosterol esters in an in vitro competitive estrogen receptor binding assay (up to 1×10^{-4} mol/L) and a recombinant yeast assay (2×10^{-4} mol/L).⁶⁷ The phytosterols tested consisted of a mixture of β -sitosterol (47.9%), campesterol (28.8%), and stigmasterol (23.3%) and were sourced from a variety of edible vegetable oil distillates (e.g., sunflower, soya bean and rapeseed oils). The esters were phytosterols esterified with fatty acids from sunflower oil. The competitive estrogen receptor binding assay used a preparation of estrogen receptors isolated from 10-week-old Wistar rat uteri and measured the concentration-dependent substitution of [2,4,5,6-³H]estradiol at the estrogen receptor.

Four phytostanol mixtures (0, 1, 10 or 100 μ mol/L) showed no estrogenic activity in human mammary adenocarcinoma (MCF-7) cells.⁵⁹ Estrogenic activity was measured as the ability to induce proliferation of these cells. Proliferation was measured by staining the cells with the protein stain sulforhodamine B and measuring optical density. The MCF-7 cells were cultured for 6 days. 17β -Estradiol was used as a positive control. The percentage of β -sitostanol in the phytostanols, derived from vegetable oil, ranged from 58% - 67%, and campestanol ranged from 29% - 32%. The phytosterol content was < 4%. Precipitation and slight cytotoxicity were observed at the highest test concentration with all mixtures. No increased cell proliferation was observable in cells treated with phytostanols. Under the conditions of this study, the phytostanol mixtures tested showed no estrogenic activity.

In Vivo

Neither WDPSE nor VODPSE administered in feed (0, 8.3%) for 4 days influenced the uterine weights of female Wistar rats (n = 10; 17-day-old) in a Teicco assay.⁵⁹ Diethylstilbestrol (5, 10, or 20 μ g/kg) in the diet was used as positive control. Uterine weight was used as an indicator of estrogenic activity. No treatment-related effects on general condition, body weight or food consumption were observed.

β -Sitosterol, stigmasterol, and their oxidation products were inactive in a 28-day mosquito fish masculinization assay at concentrations up to 100 μ g/L.⁶⁸

There were no signs of estrogenic activity for phytosterols and phytosterol esters tested in an in vivo immature rat uterotrophic assay (n = 10; up to 500 mg/kg).⁶⁷ The phytosterols tested consisted of a mixture of β -sitosterol (47.9%), campesterol (28.8%), and stigmasterol (23.3%) and were sourced from a variety of edible vegetable oil distillates (e.g. sunflower, soya bean and rapeseed oils). The phytosterol esters were prepared by esterifying these phytosterols with fatty acids from sunflower oil.

Absolute and relative uterine weights were unaffected in an immature rat uterotrophic assay of a mixture of phytosterols and phytostanols (0, 500, 1000, 2500 mg/kg) administered twice daily for 4 days when compared with the negative control.⁵⁹ The mixture of phytosterols and phytostanols used in this study was derived by solvent extraction (~40–55% β -sitosterol, 16–31% β -sitostanol, 11–15% campesterol and 2–11% campestanol; MPSS-SE) and was assessed using female, Crl:CD (SD)IGS BR VAF/Plus, 19-day-old rats (n = 10). Ethinyl estradiol was used as a positive control. Body weight gains of animals in the 2000 and 5000 mg/kg [sic] groups were reduced.

PEGs Soy Sterol Report

Dose-dependent uterotrophic effects of subcutaneously administered β -sitosterol in ovariectomized rats and its synergism with estradiol could be attributable to intrinsic estrogenic properties; the effects of β -sitosterol could be inhibited by progesterone.⁶¹

β -Sitosterol was an effective estrogen-like agonist causing vaginal cornification and uterine weight gain in adult, ovariectomized Wistar rats.⁶⁹ Subcutaneous injections of the sterol caused dose-related increases in uterine glycogen concentration after 10 days.

Progesterone treatment partially suppressed the phytosterol-induced elevation of glycogen concentration when administered in combination with the median and high phytosterol doses. β -Sitosterol also stimulated glucose-6-phosphate dehydrogenase, phosphohexose isomerase, and total lactate dehydrogenase activities.

In a related study, uterine RNA, DNA, and protein concentrations were increased by subcutaneous treatment with β -sitosterol.⁶¹

Other studies of well-characterized phytosterols and phytosterol esters demonstrated no effect in an estrogen-binding study, a recombinant yeast assay for estrogen or estrogen-like activity, or a juvenile rat uterotrophic assay for estrogen or estrogen-like activity.^{61,67,70}

Sulfates of β -sitosterol act as abortifacients in Dutch-belted rabbits (1-2 μ g/d for 16 days) via estrogenic effects. They also exhibit spermicidal effects. β -Sitosterol itself had anti-estrogenic, anti-progestational, gonadotrophic, anti-gonadotrophic, and anti-androgenic effects.^{46,71,72}

GENOTOXICITY

In multiple in vitro (up to 5000 μ g/plate) and in vivo (up to 2000 mg/kg) assays, phytosterols and phytosterol esters were negative for genotoxicity (Table 9). These tests included reverse mutation, chromosomal aberrations, gene mutation, clastogenicity, sister chromatid exchange (mice), micronucleus induction (rats and mice), and unscheduled DNA synthesis

assays (rats).⁷³⁻⁷⁶

PEGs Soy Sterol Report

Phytosterols and phytosterol esters were not genotoxic, with or without metabolic activation, in the Ames assay, a human lymphocyte chromosome damage assay, an unscheduled DNA synthesis assay, or a rat bone marrow micronucleus assay.^{61,77-83}

CARCINOGENICITY

No new published carcinogenicity studies were discovered and no unpublished data were submitted.

PEGs Soy Sterol Report

Sitosterol inhibited the tumor-promoting activity of 12-*O*-tetradecanoylphorbol-13-acetate (TPA) in the skin of female ICR mice after initiation with 7,12-dimethylbenz[*a*]anthracene (DMBA). The percent reduction in the average number of tumors at week 18 was 40% in mice given TPA, DMBA, and sitosterol. Sitosterol (100 µg/L in acetone/DMSO, 9:1) applied to the shaved backs of the mice before treatment with TPA inhibited TPA-induced epidermal ornithine decarboxylase (ODC) activity; ODC induction can represent the effects of phorbol esters with strong tumor promoting activity. Additionally, dermal inflammation caused by a single application of TPA was slightly inhibited by sitosterol and stigmasterol.^{46,84}

Male Fischer CD rats coadministered the direct-acting carcinogen *N*-methylnitrosourea (by cannulation on days 1, 4, 7, 10) and β-sitosterol (95% pure, with 4% campesterol and 1 % stigmasterol; 0.2% in feed for 28 weeks) had significantly fewer colonic tumors (benign or benign and malignant) compared to rats given the carcinogen alone after 28 weeks.⁸⁵ Of rats given the carcinogen alone, 54% had tumors. Of rats given both the carcinogen and sitosterol, 33% had tumors. The incidence of rats with malignant colonic neoplasms increased after coadministration of the phytosterols; 15% (7/48) had invasive carcinomas in the sterol plus carcinogen group compared to 7% (5/71) of rats given the carcinogen alone.

The phytosterols decreased epithelial cell proliferation of the colon in mice (0.1 % in feed) and rats (0.2% in feed after induction with *N*-methyl-*N*-nitrosourea), and were cytotoxic for human epidermoid carcinoma of the nasopharynx (> 20 µg/ml).^{86,87}

IRRITATION AND SENSITIZATION

Irritation

Dermal – Non-Human

WDPSE (2000 mg/kg) administered to the clipped skin of male albino rabbits (n = 3) for 4 h under semi-occlusion was not irritating.⁵⁹ VODPSE caused very slight erythema after 1 h of treatment, which was completely reversed within 24 h after treatment. Skin irritation/corrosion was tested using rabbits in according to OECD Test Guideline 404.

Dermal – Human

Phytosterols (100%; 1 mL; derived from pomegranate) were not irritating to scarified skin in a repeat irritation assay (n = 10).⁸⁸ The test site was scratched with a 30-gauge needle. The test material was administered to the same scarified location on the forearm, using a chamber, for 24 h for three consecutive days. The site was examined 30 min after removal and before the next treatment.

Dermal - In-Vitro

In an EpiDerm™ assay, phytosterols (100%) from three sources (derived from pomegranate, soybean, and acai) were not predicted to be dermal irritants.⁸⁹⁻⁹¹

Ocular

There was no irritation potential revealed for WDPSE and VODPSE (concentration not provided) in a chicken enucleated eye assay.⁵⁹

WDPSE and VODPSE (concentration not provided; assumed 100%) were considered minimally irritating in a Draize assay using albino rabbits (n not provided).⁵⁹ The assay was conducted in accordance with OECD Test Guideline 405. WDPSE and VODPSE (concentration not provided; assumed 100%) caused slight and slight or moderate discharge, respectively, which was reversible within 24 h after treatment.

In an EpiOcular™ assay, phytosterols (100%) from three sources (derived from pomegranate, soybean, and acai) were not predicted to be ocular irritants.⁸⁹⁻⁹¹

Sensitization

Non-Human

Neither WDPSE nor VODPSE (concentration not provided) caused signs of skin sensitization after administration to male guinea pigs (n = 10) in a maximization assay conducted in accordance with OECD Test Guideline 406.⁵⁹

Human

There were no signs of irritation or sensitization in a human repeat insult patch test (HRIPT; n = 50) of sterols (100%; 0.2 mL; 0.2 g; derived from pomegranate).⁹² The test material was heated to liquefy it, and then it was applied to an occlusive, hypoallergenic patch. The patch was applied to the infrascapular regions of the back for nine treatments. The same concentration and amount of the test substance used in the challenge phase.

None of the subjects with confirmed soy allergies (n = 29) had a positive reaction to a skin prick test of plant stanol ester.¹³ An open challenge with plant stanol ester within four weeks of the HRIPT (cumulative dose 5.55g) was negative in 26 of 33 (the original 29 + 4 more) subjects. Positive reactions consisted of itching of the throat in three participants, cutaneous symptoms in three, and loose stools in one subject. The reactions were observed after the final cumulative dose of plant stanol ester; all symptoms resolved without treatment.

A follow-up DBPCFC study with plant stanol ester performed on 6 of the subjects with positive reactions in the HRIPT had negative results. The DBPCFC with plant stanol ester in the remaining seventh subject (female) was interpreted as negative, although she reported loose stools the morning after the last challenge, which contained plant stanol ester. In view of the cumulative oil intake, a nonimmune-mediated effect may be considered. No further details were provided.¹³

Of 22 subjects that had positive reactions to a commercial soy extract in a skin prick test, 16 had a positive reaction to soy isolate and 6 to soy.¹⁹ None had a reaction to phytosterols.

SUMMARY

A total of 26 phytosterols and steryl alkanoates are described for use in cosmetics. These ingredients are sterols derived from plants, many of which are then esterified with plant-derived fatty acids. These ingredients are reported to function as skin-conditioning agents, hair conditioning agents, viscosity increasing agents, skin protectants, antioxidants, drug astringents, and fragrances.

Relevant component chemicals that are cosmetic ingredients and have been reviewed by the Panel were all found to be safe or safe with qualifications. Butyric acid, caprylic acid/capric acid, and linoleic acid/linolenic acid have not been reviewed. Octyldecanoic acid is not a cosmetic ingredient.

Phytosterols are from edible plant sources and exposure to phytosterols in food results in a much greater systemic exposure than that resulting from use in cosmetic products containing these ingredients. It was noted in the PEG soy sterol report that phytosterols and phytosterol esters are not significantly absorbed after oral exposure. Therefore, acute and repeated dose oral toxicity potential of these phytosterols was not addressed in this report and the focus is on the potential for reproduction toxicity, genotoxicity, carcinogenicity, irritation, and sensitization.

Protein content of phytosterol blends was not detectable at the detection limits of 10 - 20 µg/g. There were no IgE-binding proteins detected in multiple hydrophilic extracts of vegetable oils samples using immunoblotting or an EAST inhibition assays.

The phytosterols are used in all of the FDA's cosmetic category groups except baby products. They are used at maximum concentrations ranging from 0.000001% - 8%.

Phytosterols are approved to be added in food products at up to 5 g/L in Europe. The Western diet contains ~160 - 360 mg/d phytosterols consisting of ~80% β-sitosterol.

Less than 5% of dietary phytosterols, phytostanols, and their esters are absorbed in the gastrointestinal tract of rats and humans.

β-sitosterol (200 µg/mL in ethanol) and β-sitosterol/campesterol (50%/40%; 200 µg/mL in ethanol) were cytotoxic to mouse macrophages in vitro.

The LD₅₀ of two mixtures of phytosterol esters was reported to be > 2000 mg/kg.

There were no maternal or teratogenic effects attributed to phytosterol esters administered in the feed of rats in a two-generation study. The NOAEL was ≥ 8.1%, the highest concentration tested. There were no signs of reproductive toxicity to male and female American minks orally administered β-sitosterol up to 50 mg/kg/d for 10 months.

Subcutaneous injections of β-sitosterol at 5 mg/kg/d for 16 to 48 days reduced sperm concentrations and fertility, and decreased testis and accessory sex tissue weights in a time-dependent manner in male rats.

In multiple in vitro (up to 5000 µg/plate) and in vivo (up to 2000 mg/kg) genotoxicity assays, phytosterols and phytosterol esters were negative. These tests included reverse mutation, chromosomal aberration, gene mutation, clastogenicity, micronucleus induction, and unscheduled DNA synthesis assays.

A phytosterol mixture was not irritating to albino rabbits at 2000 mg/kg.

Two phytosterol mixtures at 2000 mg/kg were minimally irritating to albino rabbits.

Phytosterols derived from pomegranate at 100% were not irritating to scarified skin in a human repeat irritation assay.

Two phytosterol mixtures were not sensitizing to guinea pigs. Phytosterols derived from pomegranate were not sensitizing in and HRIPT at 100%. None of 29 subjects with confirmed soy allergies had a positive reaction to a skin prick test with plant stanol ester. Of 22 subjects that had positive reactions to a commercial soy extract in a skin prick test, none had a reaction to phytosterols.

There was little or no estrogenic activity detected in phytosterols using in vitro estrogen binding assays. Two phytosterol ester mixes administered in feed at 8.3% for 4 days did not affect the uterus weights of 17-day-old rats in a

Teicco assay. However, in the PEG soy sterol report, β -Sitosterol was an effective estrogen-like agonist causing vaginal cornification and uterine weight gain in adult, ovariectomized Wistar rats.

There were no signs of relevant estrogenic activities in phytosterol mixtures up to 2500 mg/kg in immature rat uterotrophic assays.

DISCUSSION

The Panel noted that phytosterols are naturally-occurring in edible plants and are consumed in a normal diet. Therefore, oral toxicity data was not considered and the Panel concentrated on the data provided for reproductive toxicity, genotoxicity, carcinogenicity, irritation, and sensitization.

Irritation assays, several human sensitization assays, and a study using guinea pigs did not demonstrate irritation or sensitization up to 100%. There were no positive reactions to phytosterols administered in a skin prick test using subjects that showed positive reactions to soy and soy isolate. No maternal or teratogenic effects were discovered from the oral administration of phytosterols. There was no evidence of genotoxicity or carcinogenicity.

The data on estrogenic activity provided little basis for concern about such effects from dermal exposure to phytosterols.

The Expert Panel expressed concern about pesticide residues and heavy metals that may be present in botanically-derived ingredients. They stressed that the cosmetics industry should continue to use current good manufacturing practices (cGMPs) to limit impurities.

The Panel discussed the issue of incidental inhalation exposure from hair grooming aids that include glycine soja (soybean) sterols in an aerosol and a pump spray up to 0.000001% and phytosterols in a face powder up to 0.05%. There were no inhalation toxicity data available. The Expert Panel believes that the sizes of a substantial majority of the particles of these ingredients, as manufactured, are larger than those in the respirable range and/or aggregate and agglomerate to form much larger particles in formulation. Thus, the adverse effects reported using high doses of respirable particles in the inhalation studies do not indicate risks posed by use in cosmetics.

The Panel noted that 95% – 99% of droplets/particles would not be respirable to any appreciable amount. Furthermore, droplets/particles deposited in the nasopharyngeal or bronchial regions of the respiratory tract present no toxicological concerns based on the chemical and biological properties of this ingredient. Coupled with the small actual exposure in the breathing zone and the concentrations at which the ingredients are used, the available information indicates that incidental inhalation would not be a significant route of exposure that might lead to local respiratory or systemic effects.

The Panel considered other data available to characterize the potential for phytosterols to cause systemic toxicity, irritation, sensitization, reproductive and developmental toxicity, and genotoxicity. They noted little or no irritation or sensitization in multiple tests of dermal and ocular exposure, the absence of maternal or teratogenic effects, negative genotoxicity in vitro and in vivo assays (including reverse mutation, chromosomal aberration, gene mutation, clastogenicity, micronucleus induction, and unscheduled DNA synthesis assays), and lack of carcinogenicity in a lifetime oral exposure study. A detailed discussion and summary of the Panel's approach to evaluating incidental inhalation exposures to ingredients in cosmetic products is available at <http://www.cir-safety.org/cir-findings>.

CONCLUSION

The CIR Expert Panel concluded that the following phytosterol ingredients are safe in the present practices of use and concentration described in this safety assessment in cosmetics:

- brassica campestris (rapeseed) sterols
- canola sterols*
- C10-40 isoalkyl acid phytosterol esters*
- dihydrophytosteryl octyldecanoate*
- euterpe oleracea sterols
- glycine soja (soybean) sterols
- persea gratissima (avocado) sterols
- phytosterols
- phytosteryl butyrate*
- phytosteryl canolate
- phytosteryl caprylate/caprinate*
- phytosteryl hydroxystearate*
- phytosteryl isostearate
- phytosteryl linoleate*
- phytosteryl linoleate/linolenate*
- phytosteryl macadamiate
- phytosteryl nonanoate*
- phytosteryl oleate
- phytosteryl rice branate
- phytosteryl ricinoleate*
- phytosteryl sunflowerseedate*
- punica granatum sterols*
- beta-sitosterol
- beta-sitosteryl acetate*
- soy sterol acetate
- tall oil sterolphytosteryl caprylate/caprinate
- phytosteryl hydroxystearate
- phytosteryl isostearate

* Were ingredients in this group not in current use to be used in the future, the expectation is that they would be used in product categories and at concentrations comparable to others in this group.

TABLES

Table 1. Definitions and functions of the phytosterols in this safety assessment.¹ Descriptions provided below in *italics* have been generated by CIR staff.

Ingredient CAS No.	Definition	Function
Brassica campestris (rapeseed) sterols	A mixture of sterols obtained from <i>Brassica campestris</i> (rapeseed) Seed Oil. <i>Rapeseed oil is known to contain brassicasterol, poriferasterol, and campesterol.</i> ⁴	Skin-conditioning agent – emollient
Canola Sterols	A mixture of sterols obtained from the seeds of the canola plant.	Skin-conditioning agent – emollient
C10-40 isoalkyl acid phytosterol esters	A complex mixture of esters of phytosterol and C10-40 isoalkyl acid.	Hair conditioning agent; skin-conditioning agent – emollient; viscosity increasing agent – nonaqueous
Dihydrophytosteryl octyldecanoate	The ester of dihydrophytosterol and branched chain octyldecanoic acid.	Skin conditioning agent – occlusive
Euterpe oleracea sterols	The sterol fraction isolated from the whole plant of <i>Euterpe oleracea</i> .	Skin conditioning agent – miscellaneous
Glycine soja (soybean) sterols	A mixture of phytosterols obtained from the soybean, <i>Glycine soja</i> . <i>Soybean is known to contain stigmasterol.</i> ⁴	Skin-conditioning agent – emollient
Persea gratissima (avocado) sterols	A mixture of sterols obtained from <i>Persea gratissima</i> (avocado) oil.	Skin-conditioning agent – emollient
Phytosterols	A mixture of sterols obtained from higher plants.	Skin conditioning agent – miscellaneous
Phytosteryl butyrate	The ester of phytosterols and butyric acid.	Hair conditioning agent; skin-conditioning agent – miscellaneous
Phytosteryl canolate	The ester of phytosterols and the fatty acids derived from canola oil.	Skin protectant; skin-conditioning agent – emollient; viscosity increasing agent – nonaqueous
Phytosteryl caprylate/caprinate	The ester of phytosterols with a mixture of caprylic acid and capric acid.	Hair conditioning agent; skin-conditioning agent – occlusive
Phytosteryl hydroxystearate	The ester of phytosterols and hydroxystearic acid.	Skin-conditioning agent – emollient
Phytosteryl isostearate	The ester of phytosterols and isostearic acid.	Hair conditioning agent; skin-conditioning agent – occlusive
Phytosteryl linoleate	The ester of phytosterols with linoleic acid.	Antioxidant
Phytosteryl linoleate/linolenate	The ester of phytosterols with a mixture of linoleic acid and linoleic acid.	Antioxidant
Phytosteryl macadamiate	The ester of phytosterols and the fatty acids derived macadamia seed oil.	Hair conditioning agent; skin-conditioning agent – miscellaneous
Phytosteryl nonanoate	The ester of phytosterols and nonanoic acid.	Hair conditioning agent; skin-conditioning agent – miscellaneous
Phytosteryl oleate	The ester of phytosterols and oleic acid.	Hair conditioning agent; skin-conditioning agent – miscellaneous
Phytosteryl rice branate	The ester of phytosterols and rice bran acid.	Drug astringent – skin protectant drug; hair conditioning agent; humectant; skin protectant; skin-conditioning agent – emollient
Phytosteryl ricinoleate	The ester of phytosterols and ricinoleic acid.	Hair conditioning agent; skin-conditioning agent – miscellaneous
Phytosteryl sunflowerseedate	The ester formed by the reaction of sunflower seed acid with phytosterols.	Skin-conditioning agent – miscellaneous
Punica granatum sterols	A mixture of sterols obtained from <i>Punica granatum</i> seed oil.	Hair conditioning agent; ; skin-conditioning agent – emollient; skin-conditioning agent – occlusive
Beta-sitosterol 83-46-5	<i>A sterol that is found in most plant oils and conforms to the structure in Figure 1.</i> ⁴	Fragrance ingredient; Skin-conditioning agent – miscellaneous
Beta-sitosteryl acetate 915-05-9	The ester of beta-sitosterol and acetic acid <i>that conforms to the structure in Figure 1.</i>	Skin-conditioning agent – miscellaneous
Soy sterol acetate	The acetic acid esters of soy sterol.	Skin-conditioning agent – occlusive
Tall oil sterol	The complex mixture of phytosterols (polycyclic polyterpenes, complex monohydric alcohols and their esters) recovered from fractions of tall oil.	Skin-conditioning agent – miscellaneous

Table 2. CIR safety assessments of constituents of phytosterol ingredients.

Constituent	Conclusion	Maximum concentration of use reported	Reference
Acetic acid	Safe as used.	0.3%	103
Brassica campestris (rapeseed)-derived ingredients	Not reviewed.	-	
Butyric acid	Not reviewed.	-	
Caprylic acid/capric acid	Not reviewed.	-	
Euterpe oleracea-derived ingredients	Not reviewed.	-	
Hydroxystearic acid	Safe as used.	10%	95
C10-40 isoalkyl acid	<i>As C10-40 isoalkyl acid octylodecanol esters, C4-5 isoalkyl cocoate, C32-36 isoalkyl stearate, and ethylhexyl C10-40 isoalkyl acidate.</i> Safe in the present practices of use and concentration described in this safety assessment when formulated to be non-irritating.	78%	94
Isostearic acid	Safe as used.	26%	96,97
Linoleate/linoleic acid	Not reviewed.	-	
Nonanoic acid	<i>As pelargonic acid.</i> Safe as used.	74%	98
Octyldecanoic acid	Not a cosmetic ingredient.	-	
Oleic acid	Safe as used.	43%	99,100
PEG-5, -10, -16, -25, -30, and -40 soy sterol	Insufficient; Safe as used.	2%	2,3
persea gratissima (avocado)-derived ingredients	Not reviewed.		
Plant-derived fatty acid oils (brassica campestris(rapeseed oil), canola oil)	Safe as used.	100%	93
Punica Granatum-derived ingredients	Not reviewed.		
Rice bran acid	Safe as used.	100%	93,101
Ricinoleic acid	Safe as used.	69%	102
Sunflower seed acid	Safe as used.	100%	93
Tall oil acid	Safe as used.	8%	104

Table 3. Chemical characterization of a single sample and multiple samples of plant sterol material (source plant not provided) demonstrating the variation in sterol content.^{3,9,10}

Phytosterol	Distribution of phytosterols (%)	
	Single sample	Five samples from five batches
Brassicasterol	1.1	2.7-3.1
Campesterol	25.8	26.5-27.0
Stigmasterol	21.6	17.4-18.1
β -Sitosterol	48.7	50.8-51.2
β -Sitostanol	1.8	Not provided
Cholesterol	0.4	0.2-0.3
Other sterols	0.8	1.2-1.7

Table 4. Percent distribution of phytosterols from common vegetable oils.^{3,5}

Oil source	Brassicasterol	Campesterol	Stigmasterol	B-Sitosterol	Δ 7 Stigmastenol	Unknown
Cocoa butter		8-11	24-31	59-62		
Coconut	2	6-9	18-19	69-75		
Corn		10-20	Trace-6	74-89		1
Cottonseed	Trace-1	8		89-91		
Linseed	2	28	10	53	4	
Olive		1-3	2	80-97		18
Palm		20-21	12-13	62-67		
Peanut	1	10-19	6-12	70-76		
Rapeseed	5-19	22-37		52-62		
Rice bran		14-33	3-6	55-63		
Safflower		8-13	4-9	52-57		23
Soybean		15-21	10-24	57-72		1
Sunflower		11-12	8-12	62-75	20	

Table 5. Chemical and physical properties of representative sterols.

Property	Value	Reference
Vegetable oil sterols		
Physical Form	Crystalline waxy powder or prills	¹¹
	Waxy, free-flowing granular powder	¹⁹
Color	White to off white	¹¹
Odor	Vegetable oil-like	¹⁹
Melting Point °C	138-158	¹¹
Water Solubility g/L @	< 0.01	¹¹
Other Solubility		¹¹
Fat at ambient temperature	2.5%	
Acetone	Soluble	
Ethyl acetate	Soluble	
Isopropanol	Soluble	
Tall oil sterols/stanols		
Physical Form	Crystalline waxy powder or prills	¹¹
Color	White to off white	¹¹
Melting Point °C	138-158	¹¹
Water Solubility g/L @ °C & pH	< 0.01	¹¹
Other Solubility		¹¹
Fat at ambient temperature	2.5%	
Acetone	Soluble	
Ethyl acetate	Soluble	
Isopropanol	Soluble	

Table 6. Comparison of phytosterol content of tall oil extracted by simpler saponification process and a more complicated, multi-step processes.⁴

Phytosterol	Saponification process (%)	Multi-step process (%)
Total phytosterols	98.1	99.7
Major phytosterols	88.7	92.7
β-Sitosterol	49.1	59.8
β-Sitostanol	19.9	23.2
Campesterol	15.0	6.5
Stigmasterol	< 1%	< 1%
Other phytosterols	9.3 (including stigmasterol)	7.0 (including stigmasterol)

Table 7. Allergens not detected in phytosterols derived from acai, soybean, and pomegranate.²⁰⁻²²

Alpha-isomethyl ionone	Amyl cinnamal	Anise alcohol
Benzyl alcohol	Benzyl benzoate	Benzyl cinnamate
Benzyl salicylate	Butylphenyl methylpropional	Cinnamal
Cinnamyl alcohol	Citral	Citronellol
Coumarin	Eugenol	Fanesol
Geraniol	Hexyl Cinnamal	Hydroxycetronellal
Hydroxymethylpentyl 3-cyclohexene carboxaldehyde	Isoeugenol	Limonene
Linalool	Methyl 2 octynoate	Evernia prunastri
Evernia furfuracea	Amylcinnamyl alcohol	

Table 8. Frequency of use according to duration and exposure of phytosterols.^{25,26}

Use type	Maximum Concentration (%)		Maximum Concentration (%)		Maximum Concentration (%)		Maximum Concentration (%)	
	Uses		Uses		Uses		Uses	
	Phytosteryl oleate		Phytosteryl rice branate		Punica granatum sterols		Beta-sitosterol	
Total/range	26	1.5-3	2	0.5-1	31	0.001-5	48	0.00007-0.06
<i>Duration of use</i>								
Leave-on	20	1.5-3	NR	0.5-1	29	0.1-5	46	0.00007-0.06
Rinse-off	6	NR	NR	NR	2	NR	2	NR
Diluted for (bath) use	NR	NR	NR	NR	NR	0.001	NR	NR
<i>Exposure type</i>								
Eye area	1	NR	1	1	3	NR	3	NR
Incidental ingestion	NR	1.5	NR	NR	14	0.1-5	1	0.00007-0.0008
Incidental Inhalation-sprays	NR	NR	NR	NR	NR	NR	4	NR
Incidental inhalation-powders	NR	NR	NR	NR	NR	NR	NR	0.0021
Dermal contact	26	3	2	0.5-1	15	0.001-0.5	47	0.0004-0.06
Deodorant (underarm)	NR	NR	NR	NR	NR	NR	NR	NR
Hair-noncoloring	NR	NR	NR	NR	2	NR	NR	NR
Hair-coloring	NR	NR	NR	NR	NR	NR	NR	NR
Nail	NR	NR	NR	NR	NR	NR	NR	NR
Mucous Membrane	NR	1.5	NR	NR	15	0.001-5	1	0.00007-0.0008
Baby	NR	NR	NR	NR	NR	NR	NR	NR
	Soy sterol acetate		Tall oil sterol					
Total/range	1	NR	7	0.0006-0.0046				
<i>Duration of use</i>								
Leave-on	1	NR	7	0.0045-0.0046				
Rinse-off	NR	NR	NR	0.0006				
Diluted for (bath) use	NR	NR	NR	NR				
<i>Exposure type</i>								
Eye area	NR	NR	NR	NR				
Incidental ingestion	NR	NR	NR	NR				
Incidental Inhalation-sprays	NR	NR	NR	NR				
Incidental inhalation-powders	NR	NR	NR	NR				
Dermal contact	1	NR	7	0.0006-0.0046				
Deodorant (underarm)	NR	NR	NR	NR				
Hair-noncoloring	NR	NR	NR	NR				
Hair-coloring	NR	NR	NR	NR				
Nail	NR	NR	NR	NR				
Mucous Membrane	NR	NR	NR	NR				
Baby	NR	NR	NR	NR				

NR = Not Reported; NS = Not Surveyed; Totals = Rinse-off + Leave-on Product Uses.

Note: Because each ingredient may be used in cosmetics with multiple exposure types, the sum of all exposure type uses may not equal the sum total uses.

Table 9. Genotoxicity assays of phytosterols.

Assay	Test material(s) (concentration)	Results	Reference
In vitro			
Reverse mutation <i>Salmonella typhimurium</i> (strains TA98, TA100, TA102)	7-ketositosterol (up to 5% in acetone/tween80, 3:1 v/v), 7 β -OH-sitosterol (up to 5%), 7 α -OH-sitosterol (up to 1%), 6 α -OH-3-keto-/6 β -OH-3-ketositosterol (ratio 4:3; up to 2.5%) and a mixture (up to 10%)	Negative with and without metabolic activation	⁷³
Reverse mutation <i>S. typhimurium</i> (TA98, TA100, TA1535 and TA1537)	Phytosterol mixture ^a (5–5000 μ g/plate)	Negative with and without metabolic activation	⁷⁶
Reverse mutation; <i>S. typhimurium</i> (TA98,	Phytosterol esters ^a (50-5000 μ g/plate)	Negative with and without	⁷⁶

Table 9. Genotoxicity assays of phytosterols.

Assay	Test material(s) (concentration)	Results	Reference
TA100, TA1535 and TA1537; <i>Escherichia coli</i> WP2 <i>uvrA</i> (pKM101)		metabolic activation	
Reverse mutation <i>S. typhimurium</i> (TA98, TA100, TA102, TA1535 and TA1537)	Phytosterol oxide concentrate from vegetable oil distillates (1.6-5000 µg/plate)	Negative with and without metabolic activation	74
Reverse mutation <i>S. typhimurium</i> (TA98, TA100, TA1535 and TA1537); <i>E. coli</i> WP2 <i>uvrA</i>	MPSS-SE ^c (104-1667 µg/plate)	Negative with and without metabolic activation	59
Reverse mutation <i>S. typhimurium</i> (TA98, TA100, TA1535, and TA1537); <i>E. coli</i> WP2 <i>uvrA</i>	MPSS-VD ^d (16-1000 µg/plate)	Negative with and without metabolic activation	59
Reverse mutation <i>S. typhimurium</i> (TA98, TA100, TA1535 and TA1537)	WDPSE ^e (62-5000 µg/plate)	Negative with and without metabolic activation	59
Reverse mutation <i>S. typhimurium</i> (TA98, TA100, TA1535 and TA1537)	VODPSE ^f (62-5000 µg/plate)	Negative with and without metabolic activation	59
Reverse mutations histidine-dependent <i>S. typhimurium</i> (TA98, TA100, TA1535, TA1537); tryptophan-dependent <i>E. coli</i> (WP2 <i>uvrA</i>)	Pomegranate sterols (50 mg/mL; 0.1 mL)	Negative with and without metabolic activation	105
Chromosomal aberration; Human peripheral blood lymphocytes	Phytosterol mixture ^a (40-160 µg/mL)	Negative with and without metabolic activation	76
Chromosomal aberration; Human peripheral blood lymphocytes	Phytosterol esters ^a (25-200 µg/mL)	Negative with and without metabolic activation	76
Chromosomal aberration; Human peripheral blood lymphocytes	Phytosterol oxide concentrate ^g (131.1-500 µg/mL)	Negative with and without metabolic activation	76
Chromosomal aberration; Human peripheral blood lymphocytes	MPSS-SE (100-1200 µg/mL)	Negative with and without metabolic activation	59
Chromosomal aberration; Human peripheral blood lymphocytes	MPSS-VD (31.3-1000 µg/mL)	Negative with and without metabolic activation	59
Chromosomal aberration; Chinese hamster ovary cells	WDPSE (up to 500 µg/ml)	Negative with and without metabolic activation	59
Chromosomal aberration; Chinese hamster ovary cells	VODPSE (up to 2000 µg/ml)	Negative with and without metabolic activation	59
Gene mutation; Mouse lymphoma L5178Y cells, <i>Tk</i> ⁺ / ₋ locus	Phytosterol esters ^a (5-80 µg/mL)	Negative with and without metabolic activation	74
Gene mutation; Mouse lymphoma L5178Y cells, <i>Tk</i> ⁺ / ₋ locus	MPSS-SE (5-167 µg/mL)	Negative with and without metabolic activation	59
Gene mutation; Mouse lymphoma L5178Y cells, <i>Tk</i> ⁺ / ₋ locus	WDPSE (20-500 µg/ml)	Negative with and without metabolic activation	59
Gene mutation; Mouse lymphoma L5178Y cells, <i>Tk</i> ⁺ / ₋ locus	VODPSE (125-3000 µg/ml)	Negative with and without metabolic activation	59
Clastogenicity (micronucleus induction); Human peripheral blood lymphocytes	Phytosterol oxide concentrate ^g (up to 625 µg/mL)	Negative with and without metabolic activation	74
In vivo			
Micronucleus induction; male rats, bone marrow	Phytosterol esters ^b (500-2000 mg/kg/d) for 2 days	Negative	76
Micronucleus induction; male and female rats, bone marrow	MPSS-SE (50, 500, 2000 mg/kg)	Negative	59
Unscheduled DNA synthesis; male rats, liver	Phytosterol esters ^b (800, 2000 mg/kg)	Negative	76
Micronucleus induction; male mice, blood	Triols (up to 9.4 mg/kg) and epoxides of a mixture of β-sitosterol and campesterol (67 mg/kg)	Negative	106
Sister chromatid exchange; male NIH mice (8 weeks old)	β-sitosterol (200, 400, 600, 1000 mg/kg)	Negative	75
Cellular proliferation kinetics; male NIH mice (8 weeks old)	β-sitosterol (200, 400, 600, 1000 mg/kg)	Negative	75
Mitotic index; male NIH mice (8 weeks old)	β-sitosterol (200, 400, 600, 1000 mg/kg)	Negative	75
Micronucleated polychromatic erythrocytes; male NIH mice (8 weeks old)	β-sitosterol (200, 400, 600, 1000 mg/kg)	Negative	75

^a Phytosterol composition: campesterol (26.7%), stigmasterol (17.7%), β-sitosterol (51%).

^b Phytosterol composition: campesterol (28.1%), stigmasterol (18.7%), β-sitosterol (45.5%)

^c MPSS-SE = Mixture of phytosterols and phytostanols derived from solvent extraction, which consisted of ~40–55% β-sitosterol, ~16–31% β-sitostanol, ~11–15% campesterol, and ~2–11% campestanol.

^d MPSS-VD = Mixture derived from vacuum distillation which consisted of ~63.5% β-sitosterol, ~21.7% β-sitostanol, ~6.5% campesterol and ~2.8% campestanol.

^e WDPSE = A wood-derived stanol mixture which consisted of ~94% β-sitostanol and ~6% campestanol.

^f VODPSE = A vegetable oil-derived mixture of phytostanol esters which consisted of ~68% β-sitostanol and ~32% campestanol.

^g Phytosterol oxide concentrate = ~30% phytosterol oxides.

REFERENCES

1. Gottschalck TE and Breslawec HP. International Cosmetic Ingredient Dictionary and Handbook. 14 ed. Washington, DC: Personal Care Products Council, 2012.
2. Andersen FA. Final report on the safety assessment of PEG-5, -10, -16, -25, -30, and -40 soy sterol. *International Journal of Toxicology*. 2000;19(Suppl. 1):29-46.
3. Final report of the amended safety assessment of PEG-5, -10, -16, -25, -30, and -40 soy sterol. *International Journal of Toxicology*. 2004;23(Suppl. 2):23-47.
4. DiSalvo RM. Phytosterols. Chapter: III, book 2. Schlossman ML. In: *The Chemistry and Manufacture of Cosmetics: Ingredients*. 3 ed. Carol Stream, IL: Allured Publishing Corporation; 2002:911-914.
5. Bailey's industrial oil and fat products. New York: John Wiley & Sons, 1979.
6. Carotech. 2013. Stelessterol™ 80%- Product information. Unpublished data submitted by Personal Care Products Council. 3 pages.
7. Lundmark L, Chun H, and Melby A. Soya sterols: Functional plant-derived ingredients for toiletries. *Soap Cosmetics Chemical Specialties*. 1976;52:33-34, 38, 40.
8. Personal Care Products Council. 8-8-2013. Information on Phytosterols Derived from Palm (*Elaeis guineensis*). Unpublished data submitted by Personal Care Products Council. 4 pages.
9. Unilever. Phytosterols (Ex Roche): Chemical characterization. 1996. Report No. Study AC960273. pp. 1-24. Unpublished data submitted by CTFA.
10. Unilever. Phytosterol Esters (EX Roche): Chemical characterization. 1996. Report No. AC960274. pp. 1-25. Unpublished data submitted by CTFA.
11. Cantrill R. Phytosterols, phytostanols and their esters: Chemical and technical assessment (CTA). Rome, Italy, Food and Agriculture Organization of the United Nations (FAO). 2008. <http://www.fao.org/fileadmin/templates/agns/pdf/jecfa/cta/69/Phytosterols.pdf>. pp. 1-13.
12. European Food Safety Authority (EFSA). Opinion of the NDA Panel related to a notification from Raisio Life Sciences on plant stanol esters produced from soybean oil sterols pursuant to Article 6 paragraph 11 of Directive 2000/13/EC. European Food Safety Authority. 2005. Report No. EFSA-Q-2004-131. pp. 1-6.
13. European Food Safety Authority (EFSA). Opinion of the Scientific Panel on Dietetic Products, Nutrition and Allergies on a request from the Commission related to a notification from Raisio Life Sciences on plant stanol esters produced from soybean oil sterols pursuant to Article 6, paragraph 11 of Directive 2000/13/EC - for permanent exemption from labelling. European Food Safety Authority. 2013. <http://www.efsa.europa.eu/en/scdocs/scdoc/571.htm>. Report No. EFSA-Q-2007-082. pp. 1-6.
14. Active Concepts. 2009. ABS Acai Sterols manufacturing flow chart. Unpublished data submitted by Personal Care Products Council.
15. Active Concepts. 2013. ABS Soybean Sterols manufacturing flow chart. Unpublished data submitted by Personal Care Products Council.
16. Active Concepts. 2010. ABS Pomegranate Sterols manufacturing flow chart. Unpublished data submitted by Personal Care Products Council.
17. Nordlee JA, Nienmann LM, Hefle SL, and Taylor SL. Determination of proteins in soybean oil from distinct processing steps. 2002 IFT Annual Meeting and Food Expo. 2002. ift.confex.com/ift/2002/techprogram/paper_13465.htm.
18. Zitouni N, Errahali Y, Metche M, Moutete M, Kanny G, Moneret-Vautrin DA, Nicolas JP, and Fremont S. Soy allergens are detected in some edible soy oils. *Journal of Allergy and Clinical Immunology*. 2001. 107:(February): pp.188
19. European Food Safety Authority (EFSA). Opinion of the Scientific Panel on Dietetic Products, Nutrition and Allergies on a request from the Commission related to a notification from Cognis, ADM and Cargill on vegetable oils-derived phytosterols and phytosterol esters fromn soybean sources pursuant to Article 6 paragraph 11 of Dircetive 2000/13/EC. European Food Safety Authority. 2007. <http://www.efsa.europa.eu/fr/efsajournal/pub/486.htm>. Report No. EFSA-Q-2006-162. pp. 1-8.
20. Active Concepts. 2006. Compositional breakdown ABS Pomegranate Sterols (Punica Granatum Sterols). Unpublished data submitted by Personal Care Products Council.
21. Active Concepts. 2013. Compositional breakdown AC Soybean Sterols (Glycine Soja (Soybean) Sterols). Unpublished data submitted by Personal Care Products Council.
22. Active Concepts. 2013. Compositional breakdown ABS Acai Sterols (Euterpe Oleracea Sterols). Unpublished data submitted by Personal Care Products Council.

23. Pachke A, Zunker K, Wigotzki M, and Steinhart H. Determination of the IgE-binding activity of soy lecithin and refined and non-refined soybean oils. *Journal of Chromatography B, Biomedical Sciences and Applications*. 2001;756(1-2):249-254.
24. General Mills, Inc. Plant sterols, phytosterols, or sitosterols which are comprised of a mixture of the common sterol components. 1979. Submitted by FDA in response to a FOIA request dated 3/15/1996.
25. Food and Drug Administration (FDA). Frequency of use of cosmetic ingredients. *FDA Database*. 2013. Washington, DC: FDA.
26. Personal Care Products Council. 3-21-2013. Concentration of use by FDA category: Plant sterols. Unpublished data submitted by Personal Care Products Council. 6 pages.
27. Bremmer HJ, Prud'homme de Lodder LCH, and van Engelen JGM. Cosmetics Fact Sheet: To assess the risks for the consumer; Updated version for ConsExpo 4. 2006. <http://www.rivm.nl/bibliotheek/rapporten/320104001.pdf>. Date Accessed 8-24-2011. Report No. RIVM 320104001/2006. pp. 1-77.
28. Bremmer HJ, Prud'homme de Lodder LCH, and van Engelen JGM. General Fact Sheet: Limiting conditions and reliability, ventilation, room size, body surface area; Updated version for ConsExpo 4. 2006. <http://www.rivm.nl/bibliotheek/rapporten/320104002.pdf>. Date Accessed 8-24-2011. Report No. RIVM 320104002/2006. pp. 1-31.
29. Johnsen MA. The Influence of Particle Size. *Spray Technology and Marketing*. 2004;24-27.
30. Rothe H, Fautz R, Gerber E, Neumann L, Rettinger K, Schuh W, and Gronewold C. Special aspects of cosmetic spray safety evaluations: Principles on inhalation risk assessment. *Toxicol Lett*. 8-28-2011;205(2):97-104.
31. Rothe H. Special aspects of cosmetic spray safety evaluation. 2011. Unpublished information presented to the 26 September CIR Expert Panel. Washington D.C.
32. Rothe H. Special aspects of powders in decorative cosmetics. 2011. Unpublished information presented at the 26 September 2011 CIR Expert Panel Meeting. Washington, DC.
33. Aguilar, F., Crebelli, R., Dusemund, B., Galtier, P., Gilbert, J., Gott, D. M., Gundert-Remy, U., König, J., Lambre, C., Leblanc, J. C., Mortensen, A., Mosesso, P., Parent-Massin, D., Rietjens, I. M. C. M., Stankovic, I., Tobback, P., Waalkens-Berendsen, I., Woutersen, R. A., Wright, M. C., Bemrah, N., Guertler, R., Larsen, J. C., and Thustos, Ch. Scientific opinion on the safety of stigmasterol-rich plant sterols as food additive. *EFSA Journal*. 2012;10(5):2659, 38.
34. European Food Safety Authority (EFSA). Opinion of the Scientific Panel on Dietetic products, nutrition and allergies [NDA] related to a Novel Food application from Forbes Medi-Tech for approval of plant sterol-containing milk-based beverages. European Food Safety Authority. 2003. <http://www.efsa.europa.eu/en/scdocs/scdoc/15.htm>. Report No. EFSA-Q-2003-075. pp. 1-12.
35. European Food Safety Authority (EFSA). Opinion of the Scientific Panel on Dietetic products nutrition and allergies [NDA] related to two scientific publications concerning aspects of serum levels of phytosterols. European Food Safety Authority. 2005. <http://www.efsa.europa.eu/en/scdocs/scdoc/211.htm>. Report No. EFSA-Q-2004-178. pp. 1-6.
36. European Food Safety Authority (EFSA). Statement on a request from the Commission related to a novel food application on rice drinks and added phytosterols by the Scientific Panel on Dietetic Products, Nutrition and Allergies (NDA). European Food Safety Authority. 2006. <http://www.efsa.europa.eu/it/efsajournal/pub/201.htm>. Report No. EFSA-Q-2005-242. pp. 1-4.
37. European Food Safety Authority (EFSA). Statement of the Scientific Panel on Dietetic Products, Nutrition and Allergies on a request from the Commission related to a novel food application on fruit juices and nectars with added phytosterols. European Food Safety Authority. 2006. <http://www.efsa.europa.eu/fr/efsajournal/pub/391.htm>. Report No. EFSA-Q-2006-057. pp. 1-4.
38. European Food Safety Authority (EFSA). Plant sterols and blood cholesterol: Scientific substantiation of a health claim related to plant sterols and lower/reduced blood cholesterol and reduced risk of (coronary) heart disease pursuant to Article 14 of Regulation (EC) No 1924/2006. European Food Safety Authority. 2008. <http://www.efsa.europa.eu/en/efsajournal/doc/781.pdf>. Report No. EFSA-Q-2008-085. pp. 1-12.
39. European Food Safety Authority (EFSA). Consumption of food and beverages with added plant sterols in the European Union. European Food Safety Authority. 2008. <http://www.efsa.europa.eu/en/efsajournal/doc/133r.pdf>. Report No. EFSA/DATEx 03. pp. 1-21.
40. European Food Safety Authority (EFSA). Danacol® and blood cholesterol: Scientific substantiation of a health claim related to a low fat fermented milk product (Danacol®) enriched with plant sterols/stanols and lowering/reducing blood cholesterol and reduced risk of (coronary) heart disease pursuant to Article 14 of Regulation (EC) No 1924/2006. European Food Safety Authority. 2009. <http://www.efsa.europa.eu/en/efsajournal/doc/1177.pdf>. Report No. EFSA-Q-2008-779. pp. 1-12.
41. European Food Safety Authority (EFSA). Scientific opinion on the substantiation of health claims related to plant sterols and plant stanols and maintenance of normal blood cholesterol concentration (ID 549, 550, 567, 713, 1234, 1235, 1466, 1634, 1984, 2909, 3140), and maintenance of normal prostate size and normal urination (ID 14, 1467, 1635) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. European Food Safety Authority. 2010. <http://www.efsa.europa.eu/en/efsajournal/doc/1813.pdf>. pp. 1-22.

42. Scientific Committee on Food (SCF). Opinion of the Scientific Committee on Food on a request for the safety assessment of the use of phytosterol esters in yellow fat spreads. European Commission Health & Consumer Protection Directorate-General. 2000. http://ec.europa.eu/food/fs/sc/scf/out56_en.pdf. Report No. SCF/CS/NF/DOS/1 Final. pp. 1-16.
43. Scientific Committee on Food (SCF). General view of the Scientific Committee on Food on the long-term effects of the intake of elevated levels of phytosterols from multiple dietary sources, with particular attention to the effects on beta-carotene (expressed on 26 September 2002). Brussels, Belgium, European Commission, Health & Consumer Protection Committees II, Scientific Co-operation and Networks. 2002. http://ec.europa.eu/comm/food/fs/sc/scf/out143_en.pdf. Report No. SCF/CS/NF/DOS/20 ADD 1 Final. pp. 1-23.
44. Scientific Committee on Food (SCF). Opinion of the Scientific Committee on Food on a report on post launch monitoring of "yellow fat spreads with added phytosterol esters" (Expressed on 26 September 2002. Brussels, Belgium, European Commission, Health & Consumer Protection Directorate-General, Directorate C - Scientific Opinions, C2 - Management of Scientific Committees II, Scientific Co-operation and Networks. 2002. http://ec.europa.eu/comm/food/fs/sc/scf/out144_en.pdf. Report No. SCF/CS/FN/DOS/21 ADD 2 Final. pp. 1-8.
45. Scientific Committee on Food (SCF). Opinion of the Scientific Committee on Food on applications for approval of a variety of plant sterol-enriched foods (expressed on 5 March 2003). Brussels, Belgium, European Commission, Health & Consumer Protection Directorate-General, Directorate C - Scientific Opinions, C2 - Management of Scientific Committees, Scientific Co-operation and Networks. 2003. http://ec.europa.eu/comm/food/fs/sc/scf/out174_en.pdf. Report No. SCF/CS/NF/DOS/15 ADD 2 Final. pp. 1-10.
46. Ling, W. H. and Jones, P. J. Dietary phytosterols: a review of metabolism, benefits and side effects. *Life Sciences*. 1995;57(3):195-206.
47. Sanders DJ, Minter HJ, Howes D, and Hepburn PA. The safety evaluation of phytoesterol esters. Part 6. The comparative absorption and tissue distribution of phytosterols in the rat. *Food and Chemical Toxicology*. 2000;38(6):485-491.
48. Hamada T, Goto H, Yamahira T, Sugawara T, Imaizumi K, and Ikeda I. Solubility in and affinity for the bile salt micelle of plant sterols are important determinants of their intestinal absorption in rats. *Lipids*. 2006;41(6):551-556.
49. Moghadasian MH. Pharmacological properties of plant sterols in vivo and in vitro observations. *Life Sciences*. 2000;67(6):1340-1346.
50. Heinemann T, Axtmann G, and von Bergmann K. Comparison of intestinal absorption of cholesterol with different plant sterols in man. *European Journal of Clinical Investigation*. 1993;23(12):827-831.
51. Sudhop T, Sahin Y, Lindenthal B, Hahn C, Luers C, Berthold HK, and von Bergmann K. Comparison of the hepatic clearances of campesterol, sitosterol, and cholesterol in healthy subjects suggest that efflux transporters controlling intestinal sterol absorption also regulate biliary secretion. *Gut*. 2002;51(6):860-863.
52. Bhattacharyya AK, Connor WE, and Sin DS. The origin of plant sterols in the skin surface lipids in humans: From diet to plasma to skin. *The Journal of Investigative Dermatology*. 1983;80(4):294-296.
53. Fricke CB, Schröder M, Poulsen M, von Bergmann K, Wester I, Ib Knudsen I, Mortensen A, and Dieter Lüjohann D. Increased plant sterol and stanol levels in brain of Watanabe rabbits fed rapeseed oil derived plant sterol or stanol esters. *British Journal of Nutrition*. 2007;98(5):890-899.
54. Adcox, C, Boyd L, Oehrl L, Allen J, and Fenner G. Comparative effects of phytosterol oxides and cholesterol oxides in cultured macrophage-derived cell lines. *Journal of Agriculture and Food Chemistry*. 2001;49(4):2090-2095.
55. Active Concepts. 2006. Summary of neutral red cytotoxicity assay for ABS Pomegranate Sterols. Unpublished data submitted by Personal Care Products Council.
56. Chiang HC, Tseng TH, Wang CJ, Chen CF, and Kan WS. Experimental antitumor agents from *Solanum indicum* L. *Anticancer Research*. 1991;11(5):1911-1918.
57. James WD, Berger TG, and Elston DM. Errors in metabolism. Chapter: 26. In: *Andrews' Diseases of the Skin: clinical Dermatology*. 11 ed. Saunders Elsevier; 2006:506-537.
58. Lee M-H, Lu K, and Patel SB. Genetic basis of sitosterolemia. *Opinions in Lipidology*. 2001;12(2):141-149.
59. Expert Committee on Food Additives (JECFA). Safety evaluation of certain food additives. Geneva, 2009. http://whqlibdoc.who.int/publications/2009/9789241660600_eng.pdf. pp. 1-642. Unpublished data submitted to the International Programme on Chemical Safety (IPCS).
60. Laraki L, Pelletier X, Mourot J, and Debry G. Effects of dietary phytosterols on liver lipids and lipid metabolism enzymes. *Annals of Nutrition and Metabolism*. 1993;37:129-133.
61. Malini T and Vanithakumari G. Effect of beta-sitosterol on uterine biochemistry: A comparative study with estradiol and progesterone. *Biochemistry & Molecular Biology International*. 1993;31(4):659-668.
62. Hepburn PA, Horner SA, and Smith M. Safety evaluation of phytosterol esters. Part 2. Subchronic 90-day oral toxicity study on phytosterol esters--A novel functional food. *Food and Chemical Toxicology*. 1999;37(5):521-532.

63. Waalkens-Berendsen, D. H., Wolterbeek, A. P. M., Wijnands, M. V. W., Richold, M., and Hepburn, P. A. Safety Evaluation of Phytosterol Esters. Part 3. Two-Generation Reproduction Study in Rats with Phytosterol Esters-a Novel Functional Food. *Food and Chemical Toxicology*. 1999;37(7):683-696.
64. Waalkens-Berendsen, D. H., Wolterbeek, A. P. M., Wijnands, M. V. W., Richold, M., and Hepburn, P. A. Safety evaluation of phytosterol esters. Part 3. Two-generation reproduction study in rats with phytosterol esters-a novel functional food. [Erratum to document cited in CA131:321734]. *Food and Chemical Toxicology*. 1999;37(7):683-696.
65. Nieminen P, Pölönen I, Ikonen K, Määttänen M, and Mustonen AM. Evaluation of reproductive safety of β -sitosterol on the American mink (*Neovison vison*). *Chemosphere*. 2008;71(3):493-499.
66. Malini T and Vanithakumari G. Antifertility effects of β -sitosterol in male albino rats. *Journal of Ethnopharmacology*. 1991;35(2):149-153.
67. Baker VA, Hepburn PA, Kennedy SJ, Jones PA, Lea, L. J., Sumpter JP, and Ashby J. Safety evaluation of phytoesters. Part 1. Assessment of oestrogenicity using a combination of in vivo and in vitro assays. *Food and Chemical Toxicology*. 1999;37(1):13-22.
68. van den Heuvel MR, Leusch FD, Taylor S, Shannon N, and McKague AB. Assessment of the reproductive-endocrine disrupting potential of chlorine dioxide oxidation products of plant sterols. *Environmental Science and Technology*. 2006;40(8):2594-2600.
69. Malini T and Vanithakumari G. Comparative study of the effects of β -sitosterol, estradiol and progesterone on selected biochemical parameters of the uterus of ovariectomised rats. *Journal of Ethnopharmacology*. 1992;36(1):51-55.
70. Zava DT, Dollbaum CM, and Blen M. Estrogen and pregestin bioactivity of foods, herbs, and spices. *Proceedings of the Society for Experimental Biology and Medicine*. 1998;217(3):369-378.
71. Burck PJ, Thakkar AL, and Zimmerman RE. Antifertility action of a sterol sulphate in the rabbit. *Journal of Reproduction and Fertility*. 1982;66(1):109-112.
72. Malini T and Vanithakumari G. Rat toxicity studies with β -sitosterol. *Journal of Ethnopharmacology*. 1990;28(2):221-234.
73. Koschutnig, Karin, Kemmo, Suvi, Lampi, Anna Maija, Piironen, Vieno, Fritz-Ton, Cornelia, and Wagner, Karl Heinz. Separation and isolation of \pm -sitosterol oxides and their non-mutagenic potential in the Salmonella microsome assay. *Food Chemistry*. 2009;118(1):133-140.
74. Lea, L. J., Hepburn, P. A., Wolfreys, A. M., and Baldrick, P. Safety evaluation of phytosterol esters. Part 8. Lack of genotoxicity and subchronic toxicity with phytosterol oxides. *Food and Chemical Toxicology*. 2004;42(5):771-783.
75. Paniagua-Pérez R, Madrigal-Bujaidar E, Reyes-Cadena S, Molina-Jasso D, Pérez Gallaga J, Silva-Miranda A, Velazco O, Hernández N, and Chamorro G. Genotoxic and cytotoxic studies of beta-sitosterol and pteropodine in mouse. *Journal of Biomedicine and Biotechnology*. 2005;2005(3):242-247.
76. Wolfreys AM and Hepburn PA. Safety evaluation of phytosterol esters. Part 7. Assessment of mutagenic activity of phytosterols, phytosterol esters and the cholesterol derivative, 4-cholesten-3-one. *Food and Chemical Toxicology*. 2002;40(4):461-470.
77. Huntingdon Life Sciences Ltd. Plant sterols: Bacterial mutation assay. 1998. Report No. KA980008. pp. 1-26. Unpublished data submitted by CTFA.
78. Huntingdon Life Sciences Ltd. Phytosterol esters: Metaphase chromosome analysis of human lymphocytes cultured in vitro. 1997. Report No. KC960257. pp. 1-34. Unpublished data submitted by CTFA.
79. Huntingdon Life Sciences Ltd. Phytosterols: Metaphase chromosome analysis of human lymphocytes cultured in vitro. 1997. Report No. KC960255. pp. 1-32. Unpublished data submitted by CTFA.
80. Huntingdon Life Sciences Ltd. Phytosterols: Bacterial mutation assay. 1996. Report No. KA960254. pp. 1-23. Unpublished data submitted by CTFA.
81. Huntingdon Life Sciences Ltd. Phytosterol esters: Bacterial mutation assay. 1996. Report No. KA960256. pp. 1-23.
82. Covance Laboratories LTD. Plant sterol ester SSE26698-02: Induction of micronuclei in the bone marrow of treated rats. 1999. Report No. KC990110. pp. 1-32. Unpublished data submitted by CRFA.
83. Covance Laboratories LTD. Plant sterol ester SSE26698-02: Measurement of unscheduled DNA synthesis in rat liver using an in vivo/in vitro procedure. 1999. Report No. KU990111. pp. 1-37. Unpublished data submitted by CTFA.
84. Yasukawa K, Takido M, Matsumoto T, Takeuchi M, and Nakagawa S. Sterol and triterpene derivatives from plants inhibit the effects of a tumor promoter, and sitosterol and betulinic acid inhibit tumor formation in mouse skin two-stage carcinogenesis. *Oncology*. 1991;48(1):72-76.
85. Raicht RF, Cohen BI, Fazzini EP, Sarwal AN, and Takahashi M. Protective effect of plant sterols against chemically induced colon tumors in rats. *Cancer Research*. 1980;40(2):403-405.

86. Deschner EE, Cohen BI, and Raicht RF. The kinetics of the protective effect of β -sitosterol against MNU-induced colonic neoplasia. *Journal of Cancer Research and Clinical Oncology*. 1982;103(1):49-54.
87. Janezic SA and Rao AV. Dose-dependent effects of dietary phytosterol on epithelial cell proliferation of the murine colon. *Food and Chemical Toxicology*. 1992;30(7):611-616.
88. Product Investigations Inc. 2006. Evaluation of the skin-irritating propensities of ABS Pomegranate Sterols on scarified skin. Unpublished data submitted by Personal Care Products Council.
89. Active Concepts. 2012. Summary of dermal and ocular irritation tests of ABS Pomegranate Sterols. Unpublished data submitted by Personal Care Products Council.
90. Active Concepts. 2013. Summary of dermal and ocular irritation tests of ABS Acai Sterols. Unpublished data submitted by Personal Care Products Council.
91. Active Concepts. 2013. Summary of dermal and ocular irritation tests of AC Soybean Sterols. Unpublished data submitted by Personal Care Products Council.
92. AMA Laboratories Inc. 2006. 50 Human subject repeat insult patch test skin irritation/sensitization evaluation (occlusive patch) ABS Pomegranate Sterols. Unpublished data submitted by Personal Care Products Council.
93. Burnett C, Fiume MM, Bergfeld WF, Belsito DV, Hill RA, Klaassen CD, Liebler DC, Marks Jr.JG, Shank RC, Slaga TJ, and Snyder PW. Plant-derived fatty acid oils as used in cosmetics. Washington, DC, Cosmetic Ingredient Review. 2011. pp. 1-100.
94. Fiume MM, Bergfeld WF, Belsito DV, Hill RA, Klaassen CD, Liebler DC, Marks Jr.JG, Shank RC, Slaga TJ, and Snyder PW. Amended safety assessment of alkyl esters as used in cosmetics. Washington, DC, Cosmetic Ingredient Review. 2013. pp. 1-83.
95. Johnson W, Bergfeld WF, Belsito DV, Hill RA, Klaassen CD, Liebler DC, Marks Jr.JG, Shank RC, and Snyder PW. Amended final report on the safety assessment of hydroxystearic acid. *International Journal of Toxicology*. 1999;18(Suppl. 1):1-10.
96. Annual review of cosmetic ingredients safety assessments - 2002/2003. *International Journal of Toxicology*. 2005;24(Suppl. 1):1-102.
97. Elder RL. Final report on the safety assessment of isostearic acid. *Journal of the American College of Toxicology*. 1983;2(7):61-74.
98. Johnson, Jr W, Heldreth B, Bergfeld WF, Belsito DV, Klaassen CD, Hill RA, Liebler DC, Ma, Shank RC, Slaga TJ, Snyder PW, and Andersen FA. Final report of the Cosmetic Ingredient Review Expert Panel on the safety assessment of pelargonic acid (nonanoic acid) and nonanoate esters. *International Journal of Toxicology*. 2011;30(Suppl. 3):228S-269S.
99. Annual review of cosmetic ingredient safety assessments - 2004/2005. *International Journal of Toxicology*. 2006;25(Suppl. 2):1-89.
100. Elder RL. Final report on the safety assessment of oleic acid, lauric acid, palmitic acid, myristic acid, and stearic acid. *Journal of the American College of Toxicology*. 1987;6(3):321-401.
101. Andersen FA. Amended final report on the safety assessment of oryza sativa (rice) bran oil, oryza sativa (rice) germ oil, rice bran acid, oryza sativa (rice) bran wax, hydrogenated rice bran wax, oryza sativa (rice) bran extract, oryza sativa (rice) extract, oryza sativa (rice) germ powder, oryza sativa (rice) starch, oryza sativa (rice) bran, hydrolyzed rice bran extract, hydrolyzed rice bran protein, hydrolyzed rice extract, and hydrolyzed rice protein. *International Journal of Toxicology*. 2006;25(Suppl. 2):91-120.
102. Andersen FA. Final report on the safety assessment of ricinus communis (castor) seed oil, hydrogenated castor oil, glyceryl ricinoleate, glyceryl ricinoleate SE, ricinoleic acid, potassium ricinoleate, sodium ricinoleate, zinc ricinoleate, cetyl ricinoleate, ethyl ricinoleate, glycol ricinoleate, isopropyl ricinoleate, methyl ricinoleate, and octyldodecyl ricinoleate. *International Journal of Toxicology*. 2013;26(Suppl. 3):31-77.
103. Heldreth B, Bergfeld WF, Belsito DV, Hill RA, Klaassen CD, Liebler DC, Marks Jr.JG, Shank RC, Slaga TJ, Snyder PW, and Andersen FA. Final report of the Cosmetic Ingredient Review Expert Panel on the safety assessment of methyl acetate. *International Journal of Toxicology*. 2012;31(Suppl. 1):112-136.
104. Robinson V, Bergfeld WF, Belsito DV, Klaassen CD, Marks Jr.JG, Shank RC, Slaga TJ, Snyder PW, and Andersen FA. Amended safety assessment of tall oil acid, sodium tallate, potassium tallate, and ammonium tallate. *International Journal of Toxicology*. 2009;28(Suppl 3):2525-2585.
105. BioScreen Testing Services Inc. 2007. Bacterial reverse mutation tests ABS Pomegranate Sterols. Unpublished data submitted by Personal Care Products Council.
106. Abramsson-Zetterberg L, Svensson M, and Johnsson L. No evidence of genotoxic effect in vivo of the phytosterol oxidation products triols and epoxides. *Toxicology Letters*. 2007;173(2):132-139.