

# Safety Assessment of *Butyrospermum parkii* (Shea)-Derived Ingredients as Used in Cosmetics

International Journal of Toxicology  
2024, Vol. 43(Supplement 1) 82S–95S  
© The Author(s) 2024  
Article reuse guidelines:  
[sagepub.com/journals-permissions](https://sagepub.com/journals-permissions)  
DOI: 10.1177/10915818231224230  
[journals.sagepub.com/home/ijt](https://journals.sagepub.com/home/ijt)



Christina L. Burnett\*, Wilma F. Bergfeld\*\*, Donald V. Belsito\*\*, Ronald A. Hill\*\*\*, Curtis D. Klaassen\*\*, Daniel C. Liebler\*\*\*, James G. Marks Jr\*\*\*\*, Ronald C. Shank\*\*\*, Thomas J. Slaga\*\*, Paul W. Snyder\*\*, Monice Fiume†, and Bart Heldreth‡

## Abstract

The Expert Panel for Cosmetic Ingredient Safety (Panel) assessed the safety of 13 *Butyrospermum parkii* (shea)-derived ingredients, which are most frequently reported to function in cosmetics as skin and hair conditioning agents. The Panel reviewed the available data to determine the safety of these ingredients. Because final product formulations may contain multiple botanicals, each containing similar constituents of concern, formulators are advised to be aware of these constituents and to avoid reaching levels that may be hazardous to consumers. Industry should use good manufacturing practices to limit impurities that could be present in botanical ingredients. The Panel concluded that these ingredients are safe in the present practices of use and concentration when formulated to be non-sensitizing.

## Keywords

Cosmetics, Safety, *Butyrospermum Parkii* (Shea) Butter, *Butyrospermum Parkii* (Shea) Butter Extract, *Butyrospermum Parkii* (Shea) Butter Unsaponifiables, *Butyrospermum Parkii* (Shea) Nut Extract, *Butyrospermum Parkii* (Shea) Nut Shell Powder, *Butyrospermum Parkii* (Shea) Oil, *Butyrospermum Parkii* (Shea) Seedcake Extract, Hydrogenated Shea Butter, Hydrogenated Shea Oil, Hydrolyzed Shea Seedcake Extract, Shea Butter Glyceride, Shea Butter Glycerides, Shea Oleine

## Introduction

The *Butyrospermum parkii* (shea)-derived ingredients detailed in this report function mainly as skin and hair conditioning agents in personal care products according to the *International Cosmetic Ingredient Dictionary and Handbook (Dictionary)*.<sup>1</sup> This report assesses the safety of the following 13 *Butyrospermum parkii* (shea)-derived ingredients:

Butyrospermum Parkii (Shea) Butter	Hydrogenated Shea Butter
Butyrospermum Parkii (Shea) Butter Extract	Hydrogenated Shea Oil
Butyrospermum Parkii (Shea) Butter Unsaponifiables	Hydrolyzed Shea Seedcake Extract
Butyrospermum Parkii (Shea) Nut Extract	Shea Butter Glyceride
Butyrospermum Parkii (Shea) Nut Shell Powder	Shea Butter Glycerides
Butyrospermum Parkii (Shea) Oil	Shea Oleine
Butyrospermum Parkii (Shea) Seedcake Extract	

The Panel previously reviewed the safety of *Butyrospermum Parkii* (Shea) Oil, *Butyrospermum Parkii* (Shea)

Butter, *Butyrospermum Parkii* (Shea) Butter Unsaponifiables, and Hydrogenated Shea Butter in the 2011 safety assessment of plant-derived fatty acid oils and found these ingredients to be safe as used in cosmetics.<sup>2</sup> Because data from the previous assessment may help to inform the safety of the ingredients listed in this current assessment, the relevant information has been summarized in this report.

Botanicals such as *Butyrospermum parkii* (shea)-derived ingredients may contain hundreds of constituents, some of which may have the potential to cause toxic effects. In this assessment, the Panel is reviewing the potential toxicity of

\*Cosmetic Ingredient Review Senior Scientific Analyst/Writer

\*\*Expert Panel for Cosmetic Ingredient Safety Member

\*\*\*Expert Panel for Cosmetic Ingredient Safety Former Member

†Cosmetic Ingredient Review Senior Director

‡Cosmetic Ingredient Review Executive Director

## Corresponding Author:

Bart Heldreth, Executive Director, Cosmetic Ingredient Review, 1620 L Street, NW, Suite 1200, Washington, DC 20036, USA.

Email: [cirinfo@cir-safety.org](mailto:cirinfo@cir-safety.org)

each of the *Butyrospermum parkii* (shea)-derived ingredients as a whole, complex mixture.

The ingredient names, according to the *Dictionary*, are written as listed above, without italics and without abbreviations. When referring to the tree from which these ingredients are derived, the standard scientific practice of using italics will be followed (e.g., *Butyrospermum parkii*). The shea tree is also known taxonomically as *Vitellaria paradoxa* and is referred to as such by many references and by the US Food and Drug Administration (FDA).

Shea oleine (“oleine” is an oleate triglyceride) is listed as a cosmetic ingredient in the FDA Voluntary Cosmetic Registration Program (VCRP) database, but it is not an ingredient listed in the *Dictionary*.<sup>1,3</sup> This chemical has been included in this report because use as a cosmetic ingredient may be demonstrated. (Triolein, the triester of glycerin and oleic acid, was previously reviewed by the Panel and was found safe as used in cosmetics).<sup>4</sup>

This safety assessment includes relevant published and unpublished data that are available for each endpoint that is evaluated. Published data are identified by conducting an exhaustive search of the world’s literature. A listing of the search engines and websites that are used and the sources that are typically explored, as well as the endpoints that the Panel typically evaluates, is provided on the Cosmetic Ingredient Review (CIR) website (<https://www.cir-safety.org/supplementaldoc/preliminary-search-engines-and-websites>; <https://www.cir-safety.org/supplementaldoc/cir-report-format-outline>). Unpublished data are provided by the cosmetics industry, as well as by other interested parties.

## Chemistry

### Definition and Plant Identification

The definitions and functions of the *Butyrospermum parkii* (shea)-derived ingredients included in this report are provided in Table 1. The raw materials for the *Butyrospermum parkii* (shea)-derived ingredients found in this report are obtained from the tree *Butyrospermum parkii*, which grows mainly in equatorial Africa.<sup>5-7</sup>

### Physical and Chemical Properties

Butyrospermum Parkii (Shea) Butter, depending on level of refinement, is an off-white or grey to yellowish-cream tallow-like solid, with a specific gravity of .918 at 15°C and a melting point of 37.8°C (reported range: 28-46°C).<sup>8-13</sup> Butyrospermum Parkii (Shea) Oil is a pale yellow liquid.<sup>14</sup> Shea Oleine is a clear yellow liquid, with a characteristic fatty odor and a density of .922-.928 at 20°C.<sup>15</sup>

### Method of Manufacture

The general description of the method of manufacturing of several *Butyrospermum parkii* (shea)-derived ingredients is described in Figure 1.<sup>16</sup>

**Butyrospermum Parkii (Shea) Butter Unsaponifiables.** Butyrospermum Parkii (Shea) Butter Unsaponifiables is obtained by molecular distillation and supercritical carbon dioxide extraction of Butyrospermum Parkii (Shea) Butter.<sup>17</sup>

**Butyrospermum Parkii (Shea) Nut Shell Powder.** According to a supplier, Butyrospermum Parkii (Shea) Nut Shell Powder is obtained by removing the shea seed from the shell and drying the shell in the sun, followed by grinding and sieving the resultant product.<sup>18</sup> The ground shell is then sterilized and examined by quality control.

**Butyrospermum Parkii (Shea) Seedcake Extract.** Butyrospermum Parkii (Shea) Seedcake Extract is produced by solubilizing the seedcake of *Butyrospermum parkii* in a mixture of water and butylene glycol (50%/50%, v/v), and then separating the soluble and insoluble phases, filtering, and sterilizing.<sup>19</sup>

### Composition/Impurities

The mean tocopherol concentrations and fatty acid compositions of *Butyrospermum parkii* (shea)-derived ingredients are provided in Table 2 and Table 3, respectively. While *Butyrospermum parkii* grows mainly in equatorial Africa, subtle differences in geographic location and climate affect the levels of the natural compounds, such as tocopherol and fatty acids, in *Butyrospermum parkii* (shea)-derived ingredients.<sup>5,6</sup>

A study of Butyrospermum Parkii (Shea) Butter (described as kernel fats; *n*-hexane extraction) from 36 samples from seven different countries found the principal triacylglycerols to be stearic-oleic-stearic (mean 31.2% of total triacylglycerols), stearic-oleic-oleic (27.7%), and oleic-oleic-oleic (10.8%).<sup>20</sup> Triterpene ester contents ranged from .5 to 6.5% and consisted of  $\alpha$ -amyrin cinnamate (mean 29.3% of total triterpene esters), butyrospermol cinnamate (14.8%),  $\alpha$ -amyrin acetate (14.1%), lupeol cinnamate (9.0%),  $\beta$ -amyrin cinnamate (7.6%), lupeol acetate (7.2%), butyrospermol acetate (5.8%), and  $\beta$ -amyrin acetate (4.9%) (Figure 2).

The same researchers identified the content and composition of triterpene alcohol fractions of the non-saponifiable lipids of Butyrospermum Parkii (Shea) Butter from 36 samples.<sup>21</sup> The shea kernels contained 30-54% fat, of which 2-12% were non-saponifiable lipids. Triterpene alcohol content in the non-saponifiable lipids was 22-72%. The triterpene alcohol fractions contained  $\alpha$ -amyrin,  $\beta$ -amyrin, lupeol, and butyrospermol with minor or trace amounts of  $\psi$ -taraxasterol, taraxasterol, parkeol, 24-methylene-24-dihydroparkeol, 24-methylenecycloartanol, dammaradienol, and 24-methylenedammarenol.

An analysis of the phenolic constituents of shea kernels by liquid chromatography-mass spectrometry (LC-MS) identified the following catechin compounds: gallic acid, catechin, epicatechin, epicatechin gallate, galocatechin, epigallocatechin, galocatechin gallate, and epigallocatechin gallate.<sup>7</sup> Quercetin and *trans*-cinnamic acid

**Table 1.** Definitions and Functions of the Ingredients in This Safety Assessment.<sup>1</sup>

Ingredient/CAS No.	Definition	Function
Butyrospermum Parkii (Shea) Butter CAS No. 91080-23-8; 194043-92-0	Butyrospermum Parkii (Shea) Butter is a fat obtained from the fruit of <i>Butyrospermum parkii</i> . The accepted scientific name for <i>Butyrospermum parkii</i> is <i>Vitellaria paradoxa</i> .	skin-conditioning agents – miscellaneous; skin-conditioning agents – occlusive; viscosity increasing agents - nonaqueous
Butyrospermum Parkii (Shea) Butter Extract CAS No. 91080-23-8	Butyrospermum Parkii (Shea) Butter Extract is the extract of Butyrospermum Parkii (Shea) Butter. The accepted scientific name for <i>Butyrospermum parkii</i> is <i>Vitellaria paradoxa</i> .	skin-conditioning agents - miscellaneous
Butyrospermum Parkii (Shea) Butter Unsaponifiables CAS No. 194043-92-0; 225234-14-0	Butyrospermum Parkii (Shea) Butter Unsaponifiables is the fraction of shea butter which is not saponified in the refining recovery of shea butter fatty acids. The accepted name for <i>Butyrospermum parkii</i> is <i>Vitellaria paradoxa</i> .	hair conditioning agents; skin-conditioning agents - miscellaneous
Butyrospermum Parkii (Shea) Nut Extract CAS No. 91080-23-8	Butyrospermum Parkii (Shea) Nut Extract is the extract of the nuts of <i>Butyrospermum parkii</i> . The accepted name for <i>Butyrospermum parkii</i> is <i>Vitellaria paradoxa</i> .	skin-conditioning agents - emollient
Butyrospermum Parkii (Shea) Nut Shell Powder CAS No. 91080-23-8	Butyrospermum Parkii (Shea) Nut Shell Powder is the powder obtained from the dried, ground nut shells of <i>Butyrospermum parkii</i> . The accepted scientific name for <i>Butyrospermum parkii</i> is <i>Vitellaria paradoxa</i> .	abrasives
Butyrospermum Parkii (Shea) Oil CAS No. 91080-23-8	Butyrospermum Parkii (Shea) Oil is the liquid fraction obtained from Butyrospermum Parkii (Shea) Butter. The accepted name for <i>Butyrospermum parkii</i> is <i>Vitellaria paradoxa</i> .	skin-conditioning agents – miscellaneous; skin-conditioning agents - occlusive
Butyrospermum Parkii (Shea) Seedcake Extract CAS No. 91080-23-8	Butyrospermum Parkii (Shea) Seedcake Extract is the extract of the seedcake of <i>Butyrospermum parkii</i> . The accepted name for <i>Butyrospermum parkii</i> is <i>Vitellaria paradoxa</i> .	skin protectants
Hydrogenated Shea Butter	Hydrogenated Shea Butter is the end product of the controlled hydrogenation of Butyrospermum Parkii (Shea) Butter.	skin-conditioning agents – occlusive; viscosity increasing agents - nonaqueous
Hydrogenated Shea Oil CAS No. 93333-83-6	Hydrogenated Shea Oil is the product obtained by the hydrogenation of Butyrospermum Parkii (Shea) Oil.	skin conditioning agents – emollient; skin-conditioning agents - occlusive
Hydrolyzed Shea Seedcake Extract	Hydrolyzed Shea Seedcake Extract is the hydrolysate of an extract of shea seedcake derived by acid, enzyme, or other method of hydrolysis.	not reported
Shea Butter Glyceride	Shea Butter Glyceride is the monoglyceride derived from Butyrospermum Parkii (Shea) Butter.	skin-conditioning agents – emollient; surfactants – emulsifying agents
Shea Butter Glycerides CAS No. 194043-92-0; 1016637-12-9	Shea Butter Glycerides are a mixture of mono-, di-, and triglycerides derived from Butyrospermum Parkii (Shea) Butter.	emulsion stabilizers; hair conditioning agents; skin-conditioning agents – miscellaneous; slip modifiers; surfactants – emulsifying agents; viscosity increasing agents - aqueous
Shea Oleine	Not in Dictionary.	Not in Dictionary.

were also identified. The mean kernel content of the catechin compounds was 4000 ppm with a range of 2100-9500 ppm.

**Butyrospermum Parkii (Shea) Butter Unsaponifiables.** Butyrospermum Parkii (Shea) Butter Unsaponifiables mainly contain terpene alcohols present in the butter in the form of cinnamic acid esters (including  $\alpha$ - and  $\beta$ -amyrin, lupeol, butyrospermol, and

cycloartenol) and phytosterols including  $\alpha$ -spinasterol,  $\Delta^7$ -stigmasterol, and stigmasterol).<sup>17</sup>

**Butyrospermum Parkii (Shea) Nut Shell Powder.** A supplier has reported that Butyrospermum Parkii (Shea) Nut Shell Powder does not contain any fragments or pieces of shea seed.<sup>18</sup> This ingredient consists of woody tissues and does not contain asbestos, free amines, antioxidants,

formaldehyde, monomers, nitrosamines, ethylene oxide, triethanolamine, 1,4-dioxane, or volatile organic compounds.

**Butyrospermum Parkii (Shea) Seedcake Extract.** According to a supplier, Butyrospermum Parkii (Shea) Seedcake Extract is composed of 2.8% of the seedcake extract, 50.0% butylene glycol, and 47.2% water.<sup>19</sup> This supplier also reported that alkaloids were less than the limit of sensitivity (.05 g/L in Dragendorff reagent), heavy metal composition included trace levels of nickel (.049 ppm) and lead (.478 ppm), and the sum of the concentrations of aflatoxins B1, B2, G1, and G2 was less than 1.0 µg/kg. The supplier also reported that neither the 26 allergenic compounds regulated by the European Union nor any pesticides were detected in this ingredient.

**Shea Oleine.** The primary component of shea oleine is the oleate triglyceride (Figure 3).

However, the sterol component of shea oleine is approximately 8% (w/w), of which approximately 97% is 4,4-

dimethyl sterols (mostly as esters of cinnamic acid), 2% is 4-demethylsterols and .5% is 4- $\alpha$ -methylsterols.<sup>16</sup>


## Use

### Cosmetic

The safety of the cosmetic ingredients included in this assessment is evaluated based on data received from the US FDA and the cosmetics industry on the expected use of these ingredients in cosmetics. Use frequencies of individual ingredients in cosmetics are collected from manufacturers and reported by cosmetic product category in FDA's VCRP database. Use concentration data are submitted by Industry in response to surveys, conducted by the Personal Care Products Council (Council), of maximum reported use concentrations by product category.

According to 2017 VCRP data, Butyrospermum Parkii (Shea) Butter has the most reported uses of the ingredients listed in this safety assessment in cosmetic products, with a total of 5447; more than three-fourths of the uses are in leave-on formulations (Table 4).<sup>3,22</sup> Butyrospermum Parkii (Shea) Butter Extract has the second greatest number of overall uses reported, with a total of 560; two-thirds of the uses are in leave-on formulations. The results of the concentration of use survey conducted in 2016 by the Council indicate Butyrospermum Parkii (Shea) Butter has the highest reported maximum concentration of use; it is used at up to 100% in moisturizers.<sup>23</sup> Butyrospermum Parkii (Shea) Oil has the second highest maximum concentrations of use, and is used at up to 11% in lipsticks.<sup>24</sup> No uses were reported for Hydrogenated Shea Oil or Hydrolyzed Shea Seedcake Extract.

In some cases, reports of uses were received from the VCRP, but no concentration of use data were provided. For example, Hydrogenated Shea Butter is reported to be used in 23 formulations, but no use concentration data



**Figure 1.** General description of the manufacturing of *Butyrospermum parkii* (shea)-derived ingredients.


**Table 2.** Mean Concentrations of Tocopherols in 102 Butyrospermum Parkii (Shea) Butter Samples by HPLC Analysis (µg/g).<sup>5</sup>

$\alpha$ -Tocopherol	$\beta$ -Tocopherol	$\gamma$ -Tocopherol	$\delta$ -Tocopherol	Total tocopherol
112	16	38	34	208


**Table 3.** Total Fatty Acid Composition of *Butyrospermum parkii* (Shea)-Derived Ingredients (%).<sup>2,6,46</sup>

Fatty Acids	Butyrospermum Parkii (Shea) Oil	Butyrospermum Parkii (Shea) Butter
Myristic (C14)	NR	.5
Palmitic (C16)	3.8-4.1	2.6-9
Stearic (C18)	41.2-56.8	25.6-50.2
Oleic (C18:1)	34.0-46.9	37.1-62.1
Linoleic (C18:2)	3.7-6.5	.6-10.8
Linolenic (C18:3)	NR	.5 max
Arachidic (C20)	1-2	0-3.5

NR-Not reported.



**Figure 2.** Triterpene esters.



**Figure 3.** Oleate triglyceride.

were provided. In other cases, no uses were reported to the VCRP, but a maximum use concentration was provided in the industry survey. For example, Shea Butter Glyceride was not reported in the VCRP database to be in use, but the industry survey indicated that it is used at concentrations up to .49%. It should be presumed that Shea Butter Glyceride is used in at least one cosmetic formulation.

Some of these ingredients may be used in products that can be incidentally ingested or come into contact with mucous membranes. For example, Butyrospermum Parkii (Shea) Oil is used in lipsticks at up to 11%.<sup>24</sup> Additionally, some of these ingredients were reported to be used in hair sprays, face powders, fragrances and body and hand sprays and could possibly be inhaled. For example, Butyrospermum Parkii (Shea) Seedcake Extract was reported to

be used in fragrance preparations at a maximum concentration of 4% and Butyrospermum Parkii (Shea) Butter was reported to be used in a face powder at 3%.<sup>23</sup> In practice, 95 to 99% of the droplets/particles released from cosmetic sprays have aerodynamic equivalent diameters >10  $\mu\text{m}$ , with propellant sprays yielding a greater fraction of droplets/particles below 10  $\mu\text{m}$  compared with pump sprays. Therefore, most droplets/particles incidentally inhaled from cosmetic sprays would be deposited in the nasopharyngeal and bronchial regions and would not be respirable (i.e., they would not enter the lungs) to any appreciable amount. Conservative estimates of inhalation exposures to respirable particles during the use of loose powder cosmetic products are 400-fold to 1000-fold less than protective regulatory and guidance limits for inert airborne respirable particles in the workplace.<sup>25-27</sup>

**Table 4.** Frequency and Concentration of Use According to Duration and Type of Exposure for Shea Ingredients.<sup>3,22-24,47</sup>

	# of Uses	Max Conc of Use (%)	# of Uses	Max Conc of Use (%)	# of Uses	Max Conc of Use (%)	# of Uses	Max Conc of Use (%)
	Butyrospermum Parkii (Shea) Butter		Butyrospermum Parkii (Shea) Butter Extract		Butyrospermum Parkii (Shea) Butter Unsaponifiables		Butyrospermum Parkii (Shea) Nut Extract	
Totals <sup>†</sup>	5447 <sup>a</sup>	0.0001-100	560	0.0000095-5	75	0.01-4.5	NR	0.00028-1
Duration of Use								
Leave-On	4146	0.001-100	384	0.0000095-5	72	0.015-4.5	NR	0.01-1
Rinse Off	1266	0.0001-10	169	0.00028-0.96	3	0.01-2	NR	0.00028-0.51
Diluted for (Bath) Use	35	0.05-3	7	0.05	NR	NR	NR	NR
Exposure Type								
Eye Area	249	0.1-8	25	0.5	39	0.16-0.5	NR	NR
Incidental Ingestion	510	0.01-9.4	39	0.075-1.9	3	0.25-2.55	NR	NR
Incidental Inhalation -Sprays	23; 1751 <sup>a</sup> ; 1124 <sup>b</sup>	0.1-0.33; 0.001-8 <sup>a</sup> ; 0.5 <sup>b</sup>	14; 132 <sup>a</sup> ; 111 <sup>b</sup>	0.001-0.025; 0.001-0.8 <sup>a</sup> ; 0.0001 <sup>b</sup>	6 <sup>a</sup> ; 7 <sup>b</sup>	0.5 <sup>a</sup>	NR	NR
Incidental Inhalation - Powders	9; 36 <sup>a</sup> ; 1124 <sup>b</sup>	3; 0.59 <sup>a</sup> ; 0.05-8 <sup>a</sup>	2; 8 <sup>a</sup> ; 111 <sup>b</sup>	0.015; 0.0000095-5 <sup>a</sup> ; 0.0001 <sup>b</sup>	4; 1 <sup>a</sup> ; 7 <sup>b</sup>	0.06	NR	NR
Dermal Contact	4625	0.0004-100	489	0.0001-5	67	0.051-4.5	NR	0.00028-1
Deodorant (underarm)	21 <sup>a</sup>	NR	1 <sup>a</sup>	0.05	NR	NR	NR	NR
Hair - Non-Coloring	268	0.0001-8	31	0.001-0.96	5	0.01-0.5	NR	0.01
Hair-Coloring	23	0.004-3.5	NR	NR	NR	NR	NR	NR
Nail	10	0.1-5	NR	0.01	NR	NR	NR	NR
Mucous Membrane	1422	0.0004-9.4	137	0.00028-1.9	3	0.051-2.55	NR	0.00028-0.51
Baby Products	46	0.005-7	10	0.1	1	4	NR	NR
	Butyrospermum Parkii (Shea) Nut Shell Powder		Butyrospermum Parkii (Shea) Oil		Butyrospermum Parkii (Shea) Seedcake Extract		Hydrogenated Shea Butter	
Totals <sup>†</sup>	2	0.00028-1	82	0.001-11	3	0.0002-5.5	23	NR
Duration of Use								
Leave-On	2	0.01-1	47	0.01-11	2	0.00023-5.5	12	NR
Rinse Off	NR	0.00028-0.5	32	0.001-2.5	1	0.00028-2	11	NR
Diluted for (Bath) Use	NR	NR	3	NR	NR	NR	NR	NR
Exposure Type								
Eye Area	NR	NR	2	0.5-8	NR	0.00023-5.5	NR	NR
Incidental Ingestion	NR	NR	1	0.5-11	NR	3	1	NR
Incidental Inhalation -Sprays	1 <sup>b</sup>	NR	33 <sup>a</sup> ; 6 <sup>b</sup>	1; 0.2 <sup>a</sup>	2 <sup>a</sup>	0.0095-4; 0.01 <sup>a</sup>	1; 5 <sup>a</sup> ; 3 <sup>b</sup>	NR
Incidental Inhalation - Powders	1 <sup>b</sup>	NR	1; 1 <sup>a</sup> ; 6 <sup>b</sup>	0.95-8 <sup>a</sup>	NR	0.0012-5 <sup>a</sup>	3 <sup>b</sup>	NR
Dermal Contact	2	0.00028-1	75	0.005-8	2	0.00023-5.5	15	NR
Deodorant (underarm)	NR	NR	NR	NR	NR	NR	NR	NR
Hair - Non-Coloring	NR	0.01	6	0.001-0.4	1	0.001-0.99	7	NR
Hair-Coloring	NR	NR	NR	NR	NR	NR	NR	NR
Nail	NR	NR	NR	0.1-2	NR	3-5	NR	NR
Mucous Membrane	NR	0.00028-0.0011	29	0.005-11	NR	0.00028-3	4	NR
Baby Products	NR	NR	1	NR	NR	5	NR	NR
	# of Uses	Max Conc of Use (%)	# of Uses	Max Conc of Use (%)	# of Uses	Max Conc of Use (%)	# of Uses	Max Conc of Use (%)
	Shea Butter Glycride		Shea Butter Glycrides		Shea Oleine			
Totals <sup>†</sup>	NR	0.49	39	0.49-6.5	3	NR		
Duration of Use								
Leave-On	NR	NR	29	6.5	3	NR		
Rinse Off	NR	0.49	10	0.49-2	NR	NR		
Diluted for (Bath) Use	NR	NR	NR	NR	NR	NR		
Exposure Type								
Eye Area	NR	NR	4	NR	NR	NR		
Incidental Ingestion	NR	NR	1	NR	NR	NR		
Incidental Inhalation -Sprays	NR	NR	13 <sup>a</sup> ; 5 <sup>b</sup>	NR	1; 1 <sup>a</sup>	NR		
Incidental Inhalation - Powders	NR	NR	5 <sup>b</sup>	NR	NR	NR		
Dermal Contact	NR	0.49	31	0.49-6.5	1	NR		
Deodorant (underarm)	NR	NR	NR	NR	NR	NR		
Hair - Non-Coloring	NR	NR	7	NR	2	NR		
Hair-Coloring	NR	NR	NR	NR	NR	NR		
Nail	NR	NR	NR	NR	NR	NR		
Mucous Membrane	NR	NR	3	2	NR	NR		
Baby Products	NR	NR	1	NR	NR	NR		

NR = No reported use

<sup>†</sup> Because each ingredient may be used in cosmetics with multiple exposure types, the sum of all exposure types may not equal the sum of total uses.<sup>a</sup> Number of uses includes VCRP entries for Vitellaria Paradoxa Nilotica (Shea) Butter.<sup>a</sup> It is possible these products may be sprays, but it is not specified whether the reported uses are sprays.<sup>b</sup> Not specified whether a powder or a spray, so this information is captured for both categories of incidental inhalation.<sup>c</sup> It is possible these products may be powders, but it is not specified whether the reported uses are powders.

The *Butyrospermum parkii* (shea)-derived ingredients described in this safety assessment are not restricted from use in any way under the rules governing cosmetic products in the European Union.<sup>28</sup>

## Non-Cosmetic

Butyrospermum Parkii (Shea) Oil (sheanut oil), from which many of the ingredients of this report are derived, is generally recognized as safe (GRAS) in the U.S. as a direct food additive (21CFR§184.1702). It is used in confections and

frostings, coatings of soft candy, and sweet sauces and toppings.

Refined sheanut oil is described as a component of a mixture of oils used as a cocoa butter substitute, as a coating agent, and in margarine and shortening in the *Food Chemicals Codex*, a compendium of internationally recognized standards published by the United States Pharmacopeia (USP) for the purity and identity of food ingredients.<sup>14</sup>

A triterpene-rich extract of *Butyrospermum parkii* has been reported to be used as a dietary supplement for the treatment of osteoarthritis.<sup>29</sup> Other studies have found that components of

shea extracts potentially have anti-inflammatory, antioxidant, and anti-tumor effects.<sup>30-33</sup>

## Toxicokinetic Studies

### Absorption, Distribution, Metabolism, and Excretion (ADME)

#### Animal

**Oral.** In an oral absorption and excretion study, groups of Colworth Wistar male rats were fed shea oleine in a semisynthetic diet.<sup>34</sup> In a low-dose experiment, groups of 24 rats received control feed, feed containing .5% shea oleine, or feed containing 5% shea oleine for 1 week, with control feed administered to all rats the week prior and the week following the exposure week. In a high-dose experiment, 2 groups of 15 male and 15 female rats received either 10 or 20% shea oleine in the feed for 3 weeks. In the first experiment, feces were collected and pooled weekly for each treatment group throughout weeks 2 and 3. In the second experiment, feces were collected and pooled for each treatment group in week 3 only. The dried fecal matter of the rats was then analyzed with thin-layer and gas-liquid chromatography for fecal lipid, total sterol, differential sterol levels, and, specifically, 4,4-dimethylsterols (the main sterol fraction (~97%) of shea oleine). Excretion of 4,4-dimethylsterols increased with the consumption of shea oleine. Apparent absorption was 27 to 52% and was estimated from the disappearance of 4,4-dimethylsterols from the feces. The majority of the 4,4-dimethylsterols was excreted unchanged.

#### Human

**Oral.** The oral absorption and excretion of shea oleine was studied in 4 male volunteers.<sup>34</sup> On day 3 of an 8 day period, the subjects consumed a single 25 g portion (approximately .4 g/kg) of shea oleine in mayonnaise. No other vegetable fats were consumed during the course of the study. Feces were collected on days 3 to 8 inclusively, freeze-dried, and weighed. The dried fecal matter was analyzed in the manner described above. Excretion of 4,4-dimethylsterols increased with the consumption of shea oleine, with a marked increase from baseline on days 4 and 5 and a return to approximate baseline on day 8. Absorption of 4,4-dimethylsterols was estimated to be 13 to 49%. The majority of the 4,4-dimethylsterols was excreted unchanged.

## Toxicological Studies

### Acute Toxicity Studies

No relevant published acute toxicity studies on *Butyrospermum parkii* (shea)-derived ingredients were identified in a literature search for these ingredients, and no unpublished data were submitted.

### Subchronic Toxicity Studies

**Shea Oleine.** In a 13-week rat feeding study, groups of 15 male and 15 female Colworth-Wistar rats received a diet containing 20% (w/w; 10 to 15 g/kg/day) shea oleine or hydrogenated shea oleine.<sup>35</sup> Additional groups of 15 male and 15 female rats were fed either 20% (w/w) palm oil, soy bean oil, or the hydrogenated equivalents. During the exposure period, body weight, food and water consumption, urine chemistry, and clinical pathology were assessed. Gross necropsy and microscopic examination of select tissues and organs were performed at study completion.

Results showed that shea oleine diets produced biological effects similar to those of palm oil and soy bean oil diets. Slightly reduced body weight gain was observed in rats fed either of the shea oleine diets when compared to diets with palm oil and soy bean oil. No significant differences in body weight gains were observed between rats fed hydrogenated shea oleine versus non-hydrogenated shea oleine. Slightly reduced cholesterol levels, increased aminotransferase levels, and lower triglyceride and alanine aminotransferase values were observed in rats fed non-hydrogenated diets, as were increased liver weights and reduced liver-lipid values. These changes were not considered to be biologically significant. Also considered biologically insignificant by the authors were raised alkaline phosphatase levels and increased food consumption in rats fed hydrogenated shea oleine. The authors concluded that all diets were well tolerated in the rats and considered none of the findings in this study to be adverse.<sup>35</sup>

### Chronic Toxicity Studies

**Butyrospermum Parkii (Shea) Oil and Shea Oleine.** See Carcinogenicity section below.

## Developmental and Reproductive Toxicity (DART) Studies

#### Oral

**Butyrospermum Parkii (Shea) Oil and Shea Oleine.** The reproductive toxicity potential of shea oleine and hydrogenated shea oleine was assessed in two dietary studies in rats.<sup>36</sup> In study 1, groups of 20 male and 20 female Colworth-Wistar rats received 7% (w/w; 3.5 g/kg/day) of either type of shea oleine in their diet for 20 weeks (breeding began at week 12 and lasted for 2 weeks). In study 2, groups of 50 male and 50 female Colworth-Wistar rats received 15% (w/w; 7.5 g/kg/day) of either type of shea oleine or Butyrospermum Parkii (Shea) Oil in their diets for 10 weeks (breeding began at week 2 and lasted for 1 week). Both studies also evaluated other commercially available materials, such as palm oil and cocoa butter. The rats received the test materials during pre-mating, mating, pregnancy and offspring weaning. Reproduction was assessed by counting the number of litters, pups born, and

pups surviving, and by measuring body weights at birth and at weaning on day 21. Skeletal evaluation using X-ray, clinical pathology and macroscopic examination were performed on F<sub>1</sub> rats. Parental animal parameters assessed were body weight, food consumption, clinical pathology, organ weights and macroscopic examination. Fatty acids and hydrocarbon levels were measured, and various tissues were evaluated in F<sub>0</sub> animals for lipogranulomata in Study 2.

Slightly decreased body weight gain, reduced cholesterol, and increased alkaline phosphatase activities were observed in rats treated with either shea oleine or hydrogenated shea oleine. No adverse effects on reproduction from any shea materials were observed in either study for any parameter. Results showed that shea oleine, hydrogenated shea oleine, and Butyrospermum Parkii (Shea) Oil were toxicologically comparable to the other commercially available materials used in this study. The authors concluded that there was no evidence of reproductive toxicity following dietary exposure to shea oleine, hydrogenated shea oleine, and Butyrospermum Parkii (Shea) Oil in rats at concentrations equating to 15% (7.5 g/kg/day).<sup>36</sup>

## Genotoxicity

### In Vitro

*Butyrospermum Parkii (Shea) Butter and Butyrospermum Parkii (Shea) Butter Unsaponifiables.* A material containing Butyrospermum Parkii (Shea) Butter (70%) and Butyrospermum Parkii (Shea) Butter Unsaponifiables (30%) was not mutagenic in an Ames test.<sup>17</sup> The material was tested at 50 to 5000 µg/plate, with and without metabolic activation. No further details were provided.

## Carcinogenicity

### Oral

*Butyrospermum Parkii (Shea) Oil and Shea Oleine.* The carcinogenic potential of shea oleine and Butyrospermum Parkii (Shea) Oil were evaluated in a dietary study in Colworth-Wistar rats for 104 weeks.<sup>16</sup> The study also evaluated palm oil. Groups of 50 male and 50 female rats received diets containing 15% (w/w; approximately equivalent to 7.5 g/kg/day) shea oleine, 15% (w/w) Butyrospermum Parkii (Shea) Oil, or 15% (w/w) palm oil. The rats were the offspring of the animals used in the reproduction study described above (study 2) and the test diets began at weaning (21 days of age). The following parameters were assessed: mortality, clinical signs of toxicity, body weight, feed intake, clinical pathology, organ weights and macroscopic and histopathological changes plus tumor type and incidence evaluation.

Final mortality rates for both sexes for shea oleine and Butyrospermum Parkii (Shea) Oil were in the range of 28 to 30% each, while the mortality rates for both sexes exposed to

palm oil was 40%. No clinical signs of toxicity were found after exposure to either shea test material. Decreased body weight gain and increased feed intake were observed in rats of both sexes fed either shea diets, while reduced cholesterol was observed in females fed the shea oleine diet. Increased alkaline phosphatase levels were observed in both sexes fed the Butyrospermum Parkii (Shea) Oil diet, but this value was only increased in females fed the shea oleine diet. Decreased heart weights and an increased incidence of pulmonary lipidosis were observed in rats of both sexes fed either shea diet. In females fed either shea diet, an increase in the number of hepatomas was observed, while in males fed shea oleine, increases in pancreatic exocrine adenomas and skin keratoacanthomas were observed. The increase in the incidence of hepatomas was thought to be related to the high fat content of the diets and were interpreted to be consistent with previous tumor data in high fat fed rats. The authors concluded that none of the findings in this study were adverse effects and that shea oleine and Butyrospermum Parkii (Shea) Oil showed no tumorigenic potential in the rat at 15% in the diet (7.5 g/kg/day).<sup>16</sup>

## Dermal Irritation and Sensitization Studies

### Irritation

Dermal irritation studies are summarized in Table 5.<sup>17,19,37-39</sup> A material containing Butyrospermum Parkii (Shea) Butter (70%) and Butyrospermum Parkii (Shea) Butter Unsaponifiables (30%) was non-irritating in an EpiSkin™ assay when tested undiluted and in a human primary cutaneous tolerance test at a 30% dilution in paraffin oil. Butyrospermum Parkii (Shea) Butter Extract at 5% in a moisturizer and Butyrospermum Parkii (Shea) Seedcake Extract at up to .14% were not irritating in human studies.

*Butyrospermum Parkii (Shea) Butter.* In an EpiSkin™ in vitro assay, 24.1% Butyrospermum Parkii (Shea) Butter in a lip wax was not an irritant.<sup>2</sup> In a rabbit study, Butyrospermum Parkii (Shea) Butter (concentration not reported) produced very slight erythema with or without edema in 2/3 rabbits exposed to the test material for 4 h in an irritation study utilizing occlusive patches. The erythema was resolved 3 or 4 days after patching. Butyrospermum Parkii (Shea) Butter did not cause primary cutaneous irritation when tested at up to 2%. No irritation to Butyrospermum Parkii (Shea) Butter was observed in human volunteers for in-use studies of lip gloss or body/hand massage oils at concentrations up to 45%.

### Sensitization

Dermal sensitization studies are summarized in Table 6.<sup>17,19,40-45</sup> A material containing Butyrospermum Parkii (Shea) Butter (70%) and Butyrospermum Parkii (Shea) Butter Unsaponifiables (30%) was negative in a direct peptide reactivity assay (DPRA)

**Table 5.** Dermal Irritation Studies for *Butyrospermum parkii* (Shea)-Derived Ingredients.

Test Article	Concentration/ Dose	Test Population	Procedure	Results	Reference
<b>In Vitro</b>					
70% Butyrospermum Parkii (Shea) Butter and 30% Butyrospermum Parkii (Shea) Butter Unsaponifiables	undiluted	N/A	EpiSkin™ assay in accordance with OECD TG 439; no further details provided	Non-irritating	17
<b>Human</b>					
Butyrospermum Parkii (Shea) Butter Extract	5% in a moisturizer	46 subjects	Single-blind, 4-week clinical use study; test material applied twice daily in place of regular moisturizer; study supervised by a dermatologist who conducted baseline, 2-week interim, and final exams	No irritation	37
Butyrospermum Parkii (Shea) Butter Extract	5% in a moisturizer	18 subjects	24-h single insult patch test; undiluted; occluded (Blenderm patch); no further details provided	No irritation	38
70% Butyrospermum Parkii (Shea) Butter and 30% Butyrospermum Parkii (Shea) Butter Unsaponifiables	30% diluted in paraffin oil	10 subjects	48-h primary cutaneous tolerance test; single patch; treated sites examined 30 min and 24 h post-patch removal; no further details provided	No irritation	17
Butyrospermum Parkii (Shea) Seedcake Extract (supplied as 2.8% in 47.2% water and 50.0% butylene glycol)	product diluted to 5%	23 subjects	48-h primary cutaneous tolerance test; single occlusive patch; no further details provided	48-h after application, one skin reaction observed on the treated site and one on the control site; mean irritation index = .04; product was considered non-irritating	19
Butyrospermum Parkii (Shea) Seedcake Extract	.006% in a SPF 50 cream	41 subjects	4-week cutaneous “in-use” test; once daily application to face, neck, and neckline; no sun exposure; study supervised by a dermatologist who conducted baseline and final exams	Good cutaneous acceptability	39

when tested undiluted, thereby indicating a lack of sensitizing potential. Butyrospermum Parkii (Shea) Butter Extract was non-sensitizing in human patch tests at up to 5% in formulation. Butyrospermum Parkii (Shea) Seedcake Extract at .14% did not produce sensitization reactions in 15 human subjects in a non-standard sensitization study where a 48-h cutaneous patch test was followed 6-8 weeks later with another 48-h patch test. Butyrospermum Parkii (Shea) Seedcake Extract was not irritating or sensitizing in human repeat insult patch tests (HRIPTs) at concentrations up to .42%.

**Butyrospermum Parkii (Shea) Butter.** Butyrospermum Parkii (Shea) Butter was not sensitizing in a guinea pig maximization study.<sup>2</sup> The induction concentration was 75% and the challenge concentrations were 20 and 50%. No sensitization was

observed in multiple HRIPTs with products containing Butyrospermum Parkii (Shea) Butter. Concentrations tested were up to 60%.

### Phototoxicity and Photosensitization

#### In Vitro

**Butyrospermum Parkii (Shea) Butter and Butyrospermum Parkii (Shea) Butter Unsaponifiables.** A material containing Butyrospermum Parkii (Shea) Butter (70%) and Butyrospermum Parkii (Shea) Butter Unsaponifiables (30%) was considered non-phototoxic in a 3T3 NRU assay when tested at .005 to 1 mg/mL.<sup>17</sup> This study was performed in accordance to Organization for Economic Co-operation and Development's (OECD) TG 432. No further details were provided.

**Table 6.** Dermal Sensitization Studies for *Butyrospermum Parkii* (Shea)-Derived Ingredients.

Test Article	Concentration/ Dose	Test Population	Procedure	Results	References
<b>In Vitro</b>					
70% Butyrospermum Parkii (Shea) Butter and 30% Butyrospermum Parkii (Shea) Butter Unsaponifiables	undiluted	N/A	Direct peptide reactivity assay (DPRA); performed in accordance with the European Centre for the Validation of Alternative Methods (ECVAM) protocol; reactivity of test material evaluated by monitoring peptide depletion following 24-h contact between test material and synthetic cysteine and lysine peptides; no further details were provided	Non-reactive and considered non-sensitizing	<a href="#">17</a>
<b>Human</b>					
Butyrospermum Parkii (Shea) Butter Extract	2% in a body lotion	28 healthy subjects	Maximization test; .05 mL test material applied neat under an occlusive dressing to a sodium lauryl sulfate (SLS) pre-treated site on the upper arm; five 48-h induction patches were followed 7-10 days later with challenge on naïve site	Not sensitizing	<a href="#">40</a>
Butyrospermum Parkii (Shea) Butter Extract	2% in a body lotion	26 healthy subjects	Maximization test; .05 mL test material applied neat under an occlusive dressing to a SLS pre-treated site on the upper arm; five 48-h induction patches were followed 7-10 days later with challenge on naïve site	Not sensitizing	<a href="#">41</a>
Butyrospermum Parkii (Shea) Butter Extract	5% in a face cream	25 healthy subjects	Maximization test; .05 mL test material applied neat under an occlusive dressing to a SLS pre-treated site on the upper arm; five 48-h induction patches were followed 7-10 days later with challenge on naïve site	Not sensitizing	<a href="#">42</a>
Butyrospermum Parkii(Shea) Butter Extract	1.7975% in a lipstick	104 subjects	HRIPT; .2 g test material applied to area 1 in <sup>2</sup> on upper back; semi-occluded	Not a dermal irritant or dermal sensitizer	<a href="#">43</a>
Butyrospermum Parkii (Shea) Seedcake Extract (supplied as 2.8% in 47.2% water and 50.0% butylene glycol)	product diluted to 5%	15 subjects	Subjects were those of the cutaneous tolerance study described above, in which 48 h patch test was considered the induction phase of the sensitization test; after a 6-8 week rest period, subjects were challenged with a single occlusive patch for 48 h; no further details provided	No reactions were observed	<a href="#">19</a>
Butyrospermum Parkii (Shea) Seedcake Extract (supplied as 2.8% in 47.2% water and 50.0% butylene glycol)	product diluted to 15%	104 subjects	HRIPT; occlusive patch; no further details provided	Not irritating or sensitizing; test score = .01	<a href="#">45</a>
Butyrospermum Parkii (Shea) Seedcake Extract	.006% in a SPF 50 cream	103 subjects	HRIPT; .02 mL test material applied over a 50 mm <sup>2</sup> area via occlusive patch (Finn Chambers on Scanpor) on back	Not irritating or sensitizing	<a href="#">44</a>

### Animal

**Butyrospermum Parkii (Shea) Butter.** Butyrospermum Parkii (Shea) Butter was not phototoxic in guinea pigs when tested at 10 and 20% in acetone.<sup>2</sup> The test sites were irradiated with UV-B light for 80 seconds followed by UV-A light for 80 min.

## Ocular Irritation Studies

### Animal

**Butyrospermum Parkii (Shea) Butter.** While mild conjunctival reactions were observed, undiluted Butyrospermum Parkii (Shea) Butter was considered not irritating when tested in the eyes of male rabbits.<sup>2</sup>

## Mucous Membrane Irritation Studies

### In Vitro

**Butyrospermum Parkii (Shea) Butter and Butyrospermum Parkii (Shea) Butter Unsaponifiables.** A balm containing 1.5% of the mixture Butyrospermum Parkii (Shea) Butter (70%) and Butyrospermum Parkii (Shea) Butter Unsaponifiables (30%) was considered non-irritating in a SkinEthic™ reconstituted mucous membrane model.<sup>17</sup> Approximately 10 µl of the balm was applied undiluted for 24 h. The negative control was phosphate buffer saline and the positive control was .1 and 3% sodium dodecyl sulfate. No further details were provided.

## Summary

The 13 *Butyrospermum parkii* (shea)-derived ingredients detailed in this report function mainly as skin and hair conditioning agents in personal care products. Studies on Butyrospermum Parkii (Shea) Butter and Butyrospermum Parkii (Shea) Butter Unsaponifiables included in the 2011 safety assessment of plant-derived fatty acid oils are not included in this summary. The original safety assessment found these ingredients to be safe as used in cosmetics.

According to 2017 VCRP data, Butyrospermum Parkii (Shea) Butter has the most reported uses of the ingredients listed in this safety assessment in cosmetic products, with a total of 5447; nearly three-fourths of the uses are in leave-on formulations. Butyrospermum Parkii (Shea) Butter Extract has the second greatest number of overall reported, with a total of 560; about two-thirds of the uses are in leave-on formulations. The results of the concentration of use survey conducted in 2016 by the Council indicate Butyrospermum Parkii (Shea) Butter has the highest reported maximum concentration of use; it is used at up to 100% in moisturizers. Butyrospermum Parkii (Shea) Oil is used at up to 11% in a lipstick. No uses were reported for Hydrogenated Shea Oil or Hydrolyzed Shea Seedcake Extract.

Butyrospermum Parkii (Shea) Oil is a GRAS direct food additive in the U.S. It is used as a cocoa butter substitute in

confections and frostings, coatings of soft candy, and sweet sauces and toppings. It is also used as a margarine or shortening. Components of shea extracts reportedly have potential anti-inflammatory, antioxidant, and anti-tumor effects.

Oral absorption and excretion studies of rats fed up to 20% shea oleine in a semisynthetic diet found excretion of 4,4-dimethylsterols increased with the consumption of shea oleine. Apparent absorption of shea oleine was 27 to 52%, as measured by 4,4-dimethylsterols. The majority of the 4,4-dimethylsterols was excreted unchanged. The findings for the absorption and excretion of approximately .4 g/kg in a single dose study of human volunteers were similar, with the absorption of shea oleine estimated to be 13 to 49%, as measured by 4,4-dimethylsterols.

In a 13-week rat feeding study, shea oleine or hydrogenated shea oleine (20% w/w, equivalent to 10-15 g/kg/day, for both test materials) did not produce adverse effects. No reproductive effects were observed in rats fed shea oleine, hydrogenated shea oleine, or Butyrospermum Parkii (Shea) Oil (up to 15% w/w, equivalent to 7.5 g/kg/day, for all test materials) for up to 20 weeks. No tumorigenic potential or adverse effects to shea oleine or Butyrospermum Parkii (Shea) Oil (15% w/w, equivalent to 7.5 g/kg/day) were observed in a carcinogenicity study in the offspring of the rats from the reproductive study.

A material containing Butyrospermum Parkii (Shea) Butter (70%) and Butyrospermum Parkii (Shea) Butter Unsaponifiables (30%) was not mutagenic in an Ames test.

A material containing Butyrospermum Parkii (Shea) Butter (70%) and Butyrospermum Parkii (Shea) Butter Unsaponifiables (30%) was non-irritating in an EpiSkin™ assay when tested undiluted and in a human primary cutaneous tolerance tested at a 30% dilution in paraffin oil. Butyrospermum Parkii (Shea) Butter Extract at 5% in a moisturizer and Butyrospermum Parkii (Shea) Seedcake Extract at up to .14% were not irritating in human irritation studies.

A material containing Butyrospermum Parkii (Shea) Butter (70%) and Butyrospermum Parkii (Shea) Butter Unsaponifiables (30%) was considered non-sensitizing in a DPRA when tested undiluted. Butyrospermum Parkii (Shea) Butter Extract was non-sensitizing in human patch tests at up to 5% in formulation. Butyrospermum Parkii (Shea) Seedcake Extract did not produce sensitization reaction in humans in a non-standard sensitization study (at .14%) or in HRIPTs (at up to .42%).

A material containing Butyrospermum Parkii (Shea) Butter (70%) and Butyrospermum Parkii (Shea) Butter Unsaponifiables (30%) was considered non-phototoxic in a 3T3 NRU assay when tested at .005 to 1 mg/mL.

A balm containing 1.5% of the mixture Butyrospermum Parkii (Shea) Butter (70%) and Butyrospermum Parkii (Shea) Butter Unsaponifiables (30%) was considered non-irritating in a SkinEthic™ reconstituted mucous membrane model.

No relevant published acute toxicity or case reports on *Butyrospermum parkii* (shea)-derived ingredients were

identified in a literature search for these ingredients and no unpublished data were submitted.

## Discussion

There are no irritation or sensitization data for *Butyrospermum Parkii* (Shea) Nut Extract and *Butyrospermum Parkii* (Shea) Nut Shell Powder, and no irritation or sensitization data for *Butyrospermum Parkii* (Shea) Seedcake Extract and *Butyrospermum Parkii* (Shea) Butter at maximum use concentrations (5.5 and 100% in leave-on products, respectively). However, human repeated patch tests for *Butyrospermum Parkii* (Shea) Seedcake Extract and *Butyrospermum Parkii* (Shea) Butter were negative at lower concentrations (.42 and 70%, respectively). Based on the Panel's clinical experience, the absence of adverse event reports, and the available negative safety test data (including negative results from DPRAs), the Panel does not expect dermal irritation or sensitization following exposure to these ingredients at reported use concentrations.

The Panel noted that, because botanical ingredients are complex mixtures, there is concern that multiple botanical ingredients in one formulation may each contribute to the final concentration of a single constituent. While there are no specific constituents of concern detailed for the *Butyrospermum parkii* (shea)-derived ingredients described in this safety assessment, not all constituents have been identified, especially for the nut shell and seedcake ingredients. Therefore, when formulating products, manufacturers should avoid reaching levels in final formulation of known botanical constituents that may cause sensitization or other adverse effects.

The Panel discussed the issue of incidental inhalation exposure from hair sprays, fragrance preparations, body and hand sprays, and face powders. There were no inhalation toxicity data available. The Panel noted that droplets/particles from spray and loose-powder cosmetic products would not be respirable to any appreciable amount. The potential for inhalation toxicity is not limited to respirable droplets/particles deposited in the lungs. In principle, inhaled droplets/particles deposited in the nasopharyngeal and thoracic regions of the respiratory tract may cause toxic effects depending on their chemical and other properties. However, coupled with the small actual exposure in the breathing zone and the concentrations at which the ingredients are used, the available information indicates that incidental inhalation would not be a significant route of exposure that might lead to local respiratory or systemic effects. A detailed discussion and summary of the Panel's approach to evaluating incidental inhalation exposures to ingredients in cosmetic products is available at <https://www.cir-safety.org/cir-findings>.

The Panel also expressed concern about pesticide residues, heavy metals, and other plant species that may be present in botanical ingredients. They stressed that the cosmetics industry should continue to use current good manufacturing practices (cGMPs) to limit impurities.

## Conclusion

The Expert Panel for Cosmetic Ingredient Safety concluded that the following 13 ingredients are safe in cosmetics in the present practices of use and concentration described in this safety assessment when formulated to be non-sensitizing.

Butyrospermum Parkii (Shea) Butter	Hydrogenated Shea Butter
Butyrospermum Parkii (Shea) Oil	Hydrogenated Shea Oil*
Butyrospermum Parkii (Shea) Butter Extractrowhead	Hydrolyzed Shea Seedcake Extract*
Butyrospermum Parkii (Shea) Butter Unsaponifiables	Shea Butter Glyceride
Butyrospermum Parkii (Shea) Nut Extract	Shea Butter Glycerides
Butyrospermum Parkii (Shea) Nut Shell Powder	Shea Oleine
Butyrospermum Parkii (Shea) Seedcake Extract	

*\*Not reported to be in current use. Were ingredients in this group not in current use to be used in the future, the expectation is that they would be used in product categories and at concentrations comparable to others in this group.*

## Author's Note

Unpublished sources cited in this report are available from the Director, Cosmetic Ingredient Review, 1620 L Street, NW, Suite 1200, Washington, DC 20036, USA.

## Author Contributions

The articles in this supplement were sponsored by the Cosmetic Ingredient Review.

## Declaration of Conflicting Interests

The author(s) declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: The articles in this supplement were sponsored by the Cosmetic Ingredient Review.

## Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The articles in this supplement were sponsored by the Cosmetic Ingredient Review. The Cosmetic Ingredient Review is financially supported by the Personal Care Products Council.

## References

1. Nikitakis J, Lange B. *International Cosmetic Ingredient Dictionary and Handbook*. 16th ed. Washington, DC: Personal Care Products Council, 2016.
2. Burnett CL, Fiume MM, Bergfeld WF, et al. Final report on plant-derived Fatty acid oils as used in cosmetics. *Int J Toxicol*. 2017;36(Suppl. 3):51S-129S.

3. Food and Drug Administration (FDA). Frequency of use of cosmetic ingredients. *FDA Database*. Washington, DC: FDA, 2017.
4. Johnson WJ, Bergfeld WF, Belsito DV, et al. Final report on the safety assessment of Trilaurin, Triarachidin, Tribehenin, Tricaprin, Tricaprylin, Trierucin, Triheptanoin, Triheptylundecanoin, Triisononanoin, Triisopalmitin, Triisostearin, Trilinolein, Trimyristin, Trioctanoin, Triolein, Tripalmitin, Tripalmitolein, Triricinolein, Tristerin, Triundecanoin, Glyceryl Triacetyl Hydroxystearate, Glyceryl Triacetyl Ricinoleate, and Glyceryl Stearate Diacetate. *Int J Toxicol*. 2001;20(Suppl 4):61-94.
5. Maranz S, Wiesman Z. Influence of climate on the tocopherol content of shea butter. *J Agric Food Chem*. 2004;52:2934-2937.
6. Maranz S, Wiesman Z, Bisgaard J, Bianchi G. Germplasm resources of *Vitellaria paradoxa* based on variations in fat compositions across the species distribution range. *Agrofor Syst*. 2004;60:71-76.
7. Maranz S, Wiesman Z, Garti N. Phenolic constituents of shea (*Vitellaria paradoxa*) kernels. *J Agric Food Chem*. 2003;51:6268-6273.
8. Salunkhe DK, Chavan JK, Adsule RN, Kadam SS. *World Oilseeds: Chemistry, Technology, and Utilization*. New York: Van Nostrand Reinhold, 1992.
9. Swern D, ed. *Bailey's Industrial Oil and Fat Products*. 4th ed. John Wiley & Sons, Inc., 1979.
10. John L. Seaton & Co., Ltd. *Seatons Refined Shea Nut Butter Specification*. John L. Seaton & Co. Limited. Unpublished data submitted by the Personal Care Products Council. 1 page, 2009.
11. John L. Seaton & Co., Ltd. *Seatons Shea Nut Butter*. John L. Seaton & Co., Limited. Unpublished data submitted by the Personal Care Products Council. 1 page, 2005.
12. Henry Lamotte Oils. Product Specification: Shea Butter, Solid. Unpublished data. 2010. Unpublished data submitted by the Personal Care Products Council on August 9, 2010. 1 page.
13. Cognis Care Chemicals. Data profile on Cetiol SB45 (Butyrospermum Parkii (Shea) Butter). Unpublished data. 2010. Unpublished data submitted by the Personal Care Products Council on August 9, 2010. 4 pages.
14. Council of Experts, United States Pharmacopeial Convention. *Food Chemicals Codex*. 8th ed. Rockville, MD: United States Pharmacopeia (USP), 2012.
15. Natural Sourcing. Material safety data sheet: Shea Olein. [https://naturalsourcing.com/downloads/msds/MSDS\\_Shea\\_Olein.pdf](https://naturalsourcing.com/downloads/msds/MSDS_Shea_Olein.pdf). Last Updated 2012.
16. Carthew P, Baldrick P, Hepburn PA. An assessment of the carcinogenic potential of shea oleine in the rat. *Food Chem Toxicol*. 2001;39:807-815.
17. Anonymous. 2016. Summary information Butyrospermum Parkii (Shea) Butter Unsaponifiables. Unpublished data submitted by Personal Care Products Council.
18. Anonymous. 2016. Summary information Butyrospermum Parkii (Shea) Nut Shell Powder. Unpublished data submitted by Personal Care Products Council.
19. Anonymous. 2017. Summary information Butyrospermum Parkii (Shea) Seedcake Extract. Unpublished data submitted by Personal Care Products Council.
20. Akihisa T, Kojima N, Katoh N, et al. Triacylglycerol and triterpene ester composition of shea nuts from seven African countries. *J Oleo Sci*. 2011;60(8):385-391.
21. Akihisa T, Kojima N, Katoh N, et al. Triterpene alcohol and fatty acid composition of shea nuts from seven African countries. *J Oleo Sci*. 2010;59(7):351-360.
22. Personal Care Products Council. 5-24-2016. Concentration of Use by FDA Product Category: Shea Butter Glyceride and Shea Butter Glycerides. Unpublished data submitted by Personal Care Products Council.
23. Personal Care Products Council. 4-18-2016. Updated Concentration of Use by FDA Product Category: Shea-Derived Ingredients. Unpublished data submitted by Personal Care Products Council.
24. Personal Care Products Council. 8-22-2016. Concentration of Use by FDA Product Category: Butyrospermum Parkii (Shea) Oil. Unpublished data submitted by Personal Care Products Council.
25. CIR Science and Support Committee of the Personal Care Products Council (CIR SSC). 11-3-2015. Cosmetic Powder Exposure. Unpublished data submitted by the Personal Care Products Council.
26. Aylott RI, Byrne GA, Middleton J, Roberts ME. Normal use levels of respirable cosmetic talc: Preliminary study. *Int J Cosmet Sci*. 1976;1(3):177-186.
27. Russell RS, Merz RD, Sherman WT, Siverston JN. The determination of respirable particles in talcum powder. *Food Cosmet Toxicol*. 1979;17(2):117-122.
28. European Union. Regulation (EC) No. 1223/2009 of the European Parliament and of the Council of 30 November 2009 on Cosmetic Products. Internet site accessed September 13, 2013; 2009. <https://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2009:342:0059:0209:en:PDF>
29. Cheras PA, Myers SP, Paul-Brent PA, Outerbridge KH, Nielsen GVL. Randomized double-blind placebo-controlled trial on the potential modes of action of SheaFlex70™ in osteoarthritis. *Phytother Res*. 2010;24:1126-1131.
30. Akihisa T, Kojima N, Kikuchi T, et al. Anti-inflammatory and chemopreventive effects of triterpene cinnamates and acetates from shea fat. *J Oleo Sci*. 2010;59(6):273-280.
31. Tagne RS, Telefo BP, Nyemb JN, et al. Anticancer and antioxidant activities of methanol extracts and fractions of some Cameroonian medicinal plants. *Asian Pac J Trop Med*. 2014;7(Suppl 1):S442-S447.
32. Zhang J, Kurita M, Shinozaki T, et al. Triterpene glycosides and other polar constituents of shea (*Vitellaria paradoxa*) kernels and their bioactivities. *Phytochemistry*. 2014;108:157-170.
33. Zhang J, Kurita M, Ebina K, et al. Melanogenesis-inhibitory activity and cancer chemopreventive effect of glucosylcucurbit acid from shea (*Vitellaria paradoxa*) kernels. *Chem Biodivers*. 2015;12:547-558.

34. Earl LK, Baldrick P, Hepburn PA. Studies to investigate the absorption and excretion of shea oleine sterols in rat and man. *Int J Toxicol*. 2002;21:353-359.
35. Earl LK, Baldrick P, Hepburn PA. A 13-week feeding study in the rat with shea oleine and hardened shea oleine. *Int J Toxicol*. 2002;21:13-22.
36. Baldrick P, Robinson JA, Hepburn PA. Reproduction studies in the rat with shea oleine and hardened shea oleine. *Food Chem Toxicol*. 2001;39:923-930.
37. Anonymous. 2009. Four week use study of a moisturizer containing 5% Butyrospermum Parkii (Shea) Butter Extract. Unpublished data submitted by Personal Care Products Council.
38. Anonymous. 2009. Clinical evaluation report: Human patch test (moisturizer containing 5% Butyrospermum Parkii (Shea) Butter Extract). Unpublished data submitted by Personal Care Products Council.
39. Institut d'Expertise Clinique. 2009. Clinical study for the appraisal of the cutaneous acceptability of a cosmetic investigational product, applied under normal conditions of use, (without sun exposure), for 4 weeks, in the Asian adult subject (SPF 50 creme containing 0.006% Butyrospermum Parkii (Shea) Seedcake Extract). Unpublished data submitted by Personal Care Products Council.
40. KGL Inc. 2009. An evaluation of the contact-sensitization potential of a topical coded product (body lotion containing 2% Butyrospermum Parkii (Shea) Butter Extract) in human skin by means of the maximization assay. (6925). Unpublished data submitted by Personal Care Products Council.
41. KGL Inc. 2009. An evaluation of the contact-sensitization potential of a topical coded product (body lotion containing 2% Butyrospermum Parkii (Shea) Butter Extract) in human skin by means of the maximization assay. (6736). Unpublished data submitted by Personal Care Products Council.
42. KGL Inc. 2009. An evaluation of the contact-sensitization potential of a topical coded product (face cream containing 5% Butyrospermum Parkii (Shea) Butter Extract) in human skin by means of the maximization assay. Unpublished data submitted by Personal Care Products Council.
43. Consumer Product Testing Co. 2014. Repeated insult patch test of lipstick containing 1.7975% Butyrospermum Parkii (Shea) Butter Extract. Unpublished data submitted by Personal Care Products Council.
44. Institut d'Expertise Clinique. 2009. Verification of the absence of sensitising potential and of the good cutaneous compatibility of a cosmetic investigational product, by repeated epicutaneous applications under occlusive patch (SPF 50 creme containing 0.006% Butyrospermum Parkii (Shea) Seedcake Extract). Unpublished data submitted by Personal Care Products Council.
45. Anonymous. 2017. Safety assessment of Butyrospermum parkii (Shea)-derived ingredients as used in cosmetics (updated information on Butyrospermum Parkii (Shea) Seedcake Extract included with memo 14). Unpublished data submitted by Personal Care Products Council.
46. Mital HC. Shea butter: cosmetic/drug applications. *Drug Cosmet Ind*. 1977;120:30-32.
47. Personal Care Products Council. 2-1-2017. Concentration of Use: shea oleine. Unpublished data submitted by Personal Care Products Council.