

Safety Assessment of Honey-Derived Ingredients as Used in Cosmetics

International Journal of Toxicology
2025, Vol. 44(Supplement 2) 5S–18S
© The Author(s) 2025
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/10915818251340381
journals.sagepub.com/home/ijt



Priya Cherian*, Wilma F. Bergfeld**, Donald V. Belsito**,
Curtis D. Klaassen**, Daniel C. Liebler***, James G. Marks Jr.***,
Lisa A. Peterson***, Ronald C. Shank***, Thomas J. Slaga***,
Paul W. Snyder**, Monice Fiume†, and Bart Heldreth††

Abstract

The Expert Panel for Cosmetic Ingredient Safety (Panel) assessed the safety of 7 honey-derived ingredients. All of these ingredients are reported to function in cosmetics as skin-conditioning agents. The Panel considered the available data relating to the safety of these ingredients in cosmetic formulations. Because impurities, particularly pesticides and endotoxins, may be present in these ingredients, formulators should continue to use good manufacturing practices to monitor and limit these possible impurities. The Panel concluded the honey-derived ingredients are safe in cosmetics in the present practices of use and concentration described in this safety assessment.

Keywords

Honey, Honey Cocoates, Honey Powder, Honey Extract, Hydrogenated Honey, Hydrolyzed Honey, Hydrolyzed Honey Protein, Cosmetic Ingredient Review, Expert Panel for Cosmetic Ingredient Safety, Safety, Cosmetics

Introduction

This is a safety assessment of the following 7 honey-derived ingredients as used in cosmetic formulations:

- Honey
- Honey Cocoates
- Honey Powder
- Honey Extract
- Hydrogenated Honey
- Hydrolyzed Honey
- Hydrolyzed Honey Protein

According to the web-based *International Cosmetic Ingredient Dictionary and Handbook* (INCI; *Dictionary*), all of these ingredients function as skin-conditioning agents.¹ Other reported functions include, but are not limited to, use as a flavoring agent, anti-acne agent, abrasive, binder, depilating agent, exfoliant, hair-conditioning agent, and nail-conditioning agent (Table 1). Use as an anti-acne agent is not considered a cosmetic function in the United States (US) and, therefore, does not fall under the purview of the Panel.

The *Dictionary* defines Honey Cocoates as a complex mixture of esters produced by the reaction of honey with coconut acid.¹ In 2017, the Panel published a safety assessment with the conclusion that coconut acid is safe in cosmetics in the present practices of use and concentration [as described in that safety assessment].² In addition, the main components

of Honey (i.e., fructose, glucose, maltose, and sucrose)³ were previously reviewed by the Panel; in 2019, a safety assessment was published with the conclusion that these component as ingredients are safe in the present practices of use and concentration [as described in that safety assessment].⁴

Some of the ingredients reviewed in this safety assessment may be consumed as food, and daily exposure from food use would result in much larger systemic exposures than those from use in cosmetic products. Although oral studies are included herein, the primary focus of this safety assessment is on the potential for effects from topical exposure to these ingredients as used in cosmetics.

It should be noted that there are multiple species of bees that produce honey; however, Honey, used as a cosmetic ingredient, has been reported to be produced by the honeybee

*Cosmetic Ingredient Review Senior Scientific Analyst/Writer

**Expert Panel for Cosmetic Ingredient Safety Member

***Expert Panel for Cosmetic Ingredient Safety Former Member

†Cosmetic Ingredient Review Senior Director

††Cosmetic Ingredient Review Executive Director

Corresponding Author:

Bart Heldreth, Executive Director, Cosmetic Ingredient Review, 555 13th St., NW, Suite 300W, Washington, DC 20004, USA.

Email: cirinfo@cir-safety.org

Table 1. INCI Names, Definitions, and Functions of the Honey Ingredients in This Safety Assessment.^{1,2}

Ingredient	Definition	Function
Honey 8028-66-8	Honey is a saccharic secretion gathered and stored by honeybees of the species, <i>Apis mellifera</i> , <i>Tetragonisca angustula</i> , <i>Scaptotrigona pectoralis</i> , or <i>Melipona becheii</i>	flavoring agent; humectant; skin-conditioning agent-humectant; solvent
Honey Cocoates	Honey Cocoates is a complex mixture of esters produced by the reaction of Honey with coconut acid [The fatty acid composition of coconut oil (from which coconut acid (CAS: 61788-47-4) is derived) is 0–1% caproic, 5–9% caprylic, 6–10% capric, 44–52% lauric, 13–19% myristic, 8–11% palmitic, 0–1% palmitoleic, 1–3% stearic, 5–8% oleic, 0–2.5% linoleic]	antiacne agent; film former; skin-conditioning agent – miscellaneous
Honey Extract 91052-95-5	Honey Extract is the extract of Honey	skin-conditioning agents-humectant; skin-conditioning agents-miscellaneous; solvents
Honey Powder	Honey Powder is the powder obtained from dehydrated, ground Honey	abrasives; binders; bulking agents; depilating agents; epilating agent; exfoliant; flavoring agent; hair conditioning agent; nail conditioning agent; skin-conditioning agent-miscellaneous
Hydrogenated Honey	Hydrogenated Honey is the end product of controlled hydrogenation of Honey	humectants; skin-conditioning agents-humectant; skin-conditioning agents-miscellaneous
Hydrolyzed Honey	Hydrolyzed Honey is the hydrolysate of Honey derived by acid, enzyme or other method of hydrolysis	skin-conditioning agents-humectant
Hydrolyzed Honey Protein	Hydrolyzed Honey Protein is the hydrolysate of honey protein derived by acid, enzyme or other method of hydrolysis	hair conditioning agents; skin-conditioning agents-miscellaneous

species *Apis mellifera*, *Tetragonisca angustula*, *Scaptotrigona pectoralis*, and *Melipona Becheii*.¹ In several studies, the honey used for testing was not produced by these species, but produced by a different honeybee species (e.g., *Apis dorsata*). Data from these studies have been included in the report as these may be helpful in drawing a conclusion of safety for this ingredient group. In most cases, information regarding the type of honey being tested (i.e., method of manufacture, floral source, species of producing bee) was not specified. However, if this information was available, it has been included in the report.

It is often not known how the substance being tested in a study compares to the cosmetic ingredient. In the report text, if it is known that the material being tested is a cosmetic ingredient, the INCI naming convention will be used (i.e., the names of cosmetic ingredients are capitalized (e.g., Honey Extract)). If it is not known that the test substance is the same as the cosmetic ingredient, the generic terminology, in all lowercase (e.g., honey extract), will be used.

This safety assessment includes relevant published and unpublished data for each endpoint that is evaluated. Published data are identified by conducting an exhaustive search of the world's literature. A list of the typical search engines and websites used, sources explored, and endpoints that Panel evaluates, is available on the Cosmetic Ingredient Review (CIR) website (<https://www.cir-safety.org/supplementaldoc/preliminary-search-engines-and-websites>; <https://www.cir-safety.org/supplementaldoc/cir-report-format-outline>).

Chemistry

Definition

According to the *Dictionary*, Honey (CAS No. 8028-66-8) is a saccharic secretion gathered and stored by honey bees of the species, *Apis mellifera*, *Tetragonisca angustula*, *Scaptotrigona pectoralis*, or *Melipona becheii*.¹ All ingredients reviewed in this report are derived from honey. The definitions of the ingredients included in this report are provided in Table 1.

Physical and Chemical Properties

Honey (CAS No. 8028-66-8) may be fluid, viscous, or solid, and ranges in color from clear to dark amber or black.⁵ Honey is acidic by nature; however, the pH and acidity levels vary depending upon botanical origin and geographic origin of the honey.⁶ The usual pH of honey ranges from 2 to 6. According to a manufacturer, a typical product with Honey Extract, prepared in water, is light to medium yellow in color, has a pH level of approximately 2.5–6.5 at 25°C, and is soluble in any proportion of water.⁷

Natural Occurrence

Honey is commercialized in most countries of the world.⁸ In the US alone, there are more than 300 types of honey, each with a unique flavor and color depending on the nectar source. Although there are many varieties of honey, the most common

types of commercialized honey include those from botanical sources such as acacia, alfalfa, avocado, blueberry, buckwheat, clover, eucalyptus, fireweed, manuka, orange blossom, sage, tupelo, and wildflower.

Honeybee Species

Apis mellifera, also known as the Western honey bee, is the most common honeybee species worldwide.⁹ This species was historically present across sub-Saharan Africa, Europe, parts of Western Asia, and the Middle East, and has now migrated westward to many countries, including the US. Honeybees of the *Tetragonisca angustula* species are stingless honeybees that are widely distributed in the neotropics, from Mexico to Northern Argentina.¹⁰ The *Scaptotrigona pectoralis* and *Melipona becheii* species are both stingless honeybee species found in South America.^{11,12}

Honey Production and Extraction

To produce honey, forager bees collect sugar-rich nectar from plant sources.^{13,14} Once brought back to the hive, the nectar is distributed, ingested, and regurgitated multiple times. This process involves the physiochemical transformations of nectar, during which sucrose is inverted into dextrose and fructose by enzymes originating from the hypopharyngeal glands of the bees. The regurgitation process also aids in the process of dehydration of the solution. The altered nectar solution is then spread over an empty comb. Further dehydration occurs by the draft created by the flapping of bee wings in the hive. Once approximately 80% of the water content is evaporated, the honeycomb cells are capped with wax for preservation.

Traditionally, honey is collected by first introducing smoke into the beehive to sedate or remove bees.¹⁴ The combs are then removed and squeezed to drain honey. Honey can also be extracted by placing combs in a metallic bowl containing a drainage hole. Burning embers are placed on top of the comb, and melted honey is drained and collected. In order to mechanically extract honey, caps are removed from combs, and placed in an extractor where centrifugation is performed. The honey is then sieved and collected.

Method of Manufacture

Information on the manufacture of Honey Extract and of a tradename mixture containing Honey Extract was provided by suppliers. The methods below regarding Honey Powder and honey protein are general to the processing of these ingredients, and it is unknown if they apply to cosmetic ingredient manufacture.

Honey Extract. According to one supplier, to produce Honey Extract, the honey is first extracted with a specified eluent under appropriate temperature conditions to yield a

concentrate.⁷ Typical eluents include water, butylene glycol, glycerin, and propylene glycol. The concentrate containing the phytochemical constituents is then blended with the desired diluent and preservation system to produce the final ingredient.

The manufacturing process of a tradename mixture containing 10.6% Honey Extract, 82.9% water, 4.4% propylene glycol dicaprylate/dicaprate, 1.5% phenoxyethanol, 0.3% xanthan gum, and 0.3% potassium sorbate was reported.¹⁵ A mixture of demineralized water, propylene glycol dicaprylate/dicaprate, and honey is combined with xanthan gum to create the final product. The manufacturing process of a different tradename mixture containing 16.5% Honey, 27.6% water, and 55.9% propylene glycol was also reported. Honey is extracted by a mixture of propylene glycol and water.¹⁶ The resulting product is then filtered.

Honey Powder. A honey powder, for food use, is produced by the combination of honey, an emulsifier, an anti-caking agent, and filler materials of high molecular weight that increase the glass transition temperature.¹⁷ Filler materials include starch, carboxymethyl cellulose, gum Arabic, maltodextrin, and gelatin. The mixture is then powdered by using either a spray or vacuum drying method with a filler to honey ratio of 50:50.

Honey Protein. Honey proteins can be extracted via physical and chemical methods.¹⁸ When physically extracting proteins, honey undergoes ultrafiltration and ultracentrifugation to isolate amylase before purification by ion exchange chromatography. A dialysis method can also be used to remove low molecular weight and interfering compounds by passive diffusion through a semipermeable membrane. Another physical extraction method involves the absorption of honey proteins by beads with specific properties. Combinatorial hexapeptide ligand library and C18 beads are used to capture honey peptidome from honey samples of chestnut, sunflower, eucalyptus, orange, and acacia. The honey peptidome is then filtered and eluted from the beads using a solvent system. Microwave-assisted hydrolysis is another method used to extract proteins from honey. Chemical methods to extract honey involve co-precipitation using compatible precipitants, such as a sodium tungstate solution, trichloroacetic acid, sulfosalicylic acid, or ammonium sulfate.

Composition

Honey. Honey is a mixture of carbohydrates, proteins, enzymes, amino acids, vitamins, minerals, antioxidants, and other compounds.¹⁴ Enzymes in honey include invertase, glucose oxidase, catalase, and acid phosphorylase. The sugar composition of honey is dependent upon the content of saccharides in the nectar used to produce the honey.⁵ Generally, fructose and glucose are found in honey in similar amounts, with D-fructose as the prevalent sugar. Non-saccharide honey components include proteins, free amino

Table 2. Chemical Composition of Honey.³

Constituent	g per 100 g honey
Water	17.1
Carbohydrates	82.4
Fructose	38.5
Glucose	31
Maltose	7.2
Sucrose	1.5
Proteins, amino acids, vitamins, and minerals	0.5
Calcium	0.0044–0.0092
Potassium	0.0132–0.0168
Copper	0.000003–0.0001
Iron	0.00006–0.0015
Magnesium	0.0012–0.0035
Manganese	0.00002–0.0004
Phosphorous	0.0019–0.0063
Sodium	0–0.0076
Zinc	0.00003–0.0004
Ascorbic acid	0.002–0.0024
Thiamin	<0.000006
Riboflavin	<0.00006
Niacin	<0.00036
Pantothenic acid	<0.00011
Pyridoxine (B6)	<0.00032

acids (including proline), carboxylic acids (gluconic, citric, lactic, malic, succinic, butyric, propionic), essential oils, dyes, and vitamins. An overview of a chemical composition of honey can be found in [Table 2](#).

Twenty-six amino acids have been reported in honey samples.¹⁸ Proline is the most predominant amino acid in floral honey, followed by phenylalanine and glutamic acid. Amino acids account for approximately 0.3–1% of total honey by weight.

Phenolic acids and flavonoids are also present in honey.³ The most common phenolic acids found in honey are 4-dimethylaminobenzoic acid, caffeic acid, *p*-coumaric acid, gallic acid, vallinic acid, syringic acid, and chlorogenic acid. Common flavonoids in honey include apigenin, genistein, pinocembrin, tricetin, chrysin, luteolin, quercetin, kaemferol, galangin, pinobanksin, and myricetin. The amounts of polyphenols in different honeys were quantified via high-performance liquid chromatography with diode-array detection (HPLC-DAD; [Table 3](#)). Generally, the quantity of a given polyphenol in the honey was approximately 0.2 mg/100 g honey, except for chestnut honey, which contained approximately 3 mg *p*-coumaric acid/100 g honey.¹⁹

Depending on the floral source, plant toxins may be transferred to the honey that is produced from their nectar, including secondary metabolites such as pyrrolizidine alkaloids, grayanotoxins, hyoscyamine, hyoscyne, saponin, strychnine, gelsemine, tutin, hyenanchin, oleandrin, and

oleandrogenin.²⁰ Honey collected from plants of the *Ericaceae* family (*Andromeda* sp., *Rhododendron ponticum*, *Kalmia* sp., *Lleucothoe* sp. *Lynoa* sp., *Pieris* sp.) has been shown to contain some of these toxins. Honey collected in areas where opium poppy cultivation is widespread has been reported to have narcotic effects.

Allergens, such as pollen, may also be present in honey.²¹ Ten grams of honey contains approximately 20 to 100,000 grains of pollen, which retain their allergenic properties during the honey-making process. Other allergens include secretions of pharyngeal and salivary glands of honeybee heads, and honeybee venom.

Several studies included in this report involve the use of tualang honey, which is a Malaysian multi-floral jungle honey produced by *Apis dorsata*. A comparison of the physicochemical characteristics of tualang and manuka honey (a mono-floral honey formed by *Apis mellifera*; found in New Zealand and Australia) is provided in [Table 4](#).²²

Honey Extract. The phenolic content of acacia, chestnut, orange tree, and woodland honey extracts were evaluated by HPLC.²³ All honey extract samples had similar, but quantitatively different, phenolic profiles. The woodland honey extract was richer in polyphenols compared to the other three extracts, showing high levels of caffeic acid, coumaric acid, ferulic acid, iso-ferulic acid, pinobanksin, and pinocembrin.

Impurities

Environmental contaminants of honey include heavy metals (e.g., lead, cadmium, and mercury), radioactive isotopes, organic pollutants, polychlorinated biphenyls, pesticides (insecticides, fungicides, herbicides, and bactericides), pathogenic bacteria, and genetically modified organisms.²⁴ Bee-keeping contaminants include acaricides (i.e., lipophilic synthetic compounds and nontoxic substances such as organic acids and components of essential oils), antibiotics (e.g., tetracyclines, streptomycin, sulfonamides, and chloramphenicol), and paradichlorobenzene.

A compound that is not naturally present in honey, 5-hydroxymethylfurfural (HMF), may be formed during the heating (via the Maillard reaction) or preservation (e.g., via acid-catalyzed dehydration of hexoses) of honey.^{20,25} HMF is a compound that may be mutagenic, carcinogenic, and cytotoxic. The *Codex Alimentarius* has established that the HMF concentration in honey should be lower than 80 mg/kg; however, the European Union recommends a lower limit of 40 mg/kg.²⁶

Honey Extract. According to one supplier, heavy metal testing was conducted on Honey Extract in a glycerin and water base.⁷ No antimony, arsenic, cadmium, chromium, iron, lead, mercury, or nickel was detected. In addition, no residual pesticides were detected.

Table 3. Honey Polyphenols Quantified With the HPLC-DAD Method.¹⁹

Honey	Polyphenol	Mean amount (mg per 100 g honey)
Acacia	<i>p</i> -coumaric acid	0.077 ± 0.003
Chestnut	<i>p</i> -coumaric acid	2.952 ± 0.004
Eucalyptus	quercetin	0.164 ± 0.007
Sunflower	caffeic acid	0.242 ± 0.001
Sunflower	<i>p</i> -coumaric acid	0.107 ± 0
Sunflower	kaempferol	0.205 ± 0.003
Sunflower	chrysin	0.217 ± 0.002
Thyme	<i>p</i> -coumaric acid	0.070 ± 0
Wild carrot	<i>p</i> -coumaric acid	0.223 ± 0.001

Table 4. Physicochemical Properties and Constituents of Tualang vs Manuka Honey.²²

Property	Tualang honey	Manuka honey
Appearance	Dark brown	Light to dark brown
pH	3.55–4	3.2–4.21
Moisture content	23.3%	18.7%
Total reducing sugars	67.5%	76%
Fructose	29.6%	40%
Glucose	30%	36.2%
Sucrose	0.6%	2.8%
Maltose	7.9%	1.2%
Potassium	0.51%	1%
Calcium	0.18%	1%
Magnesium	0.11%	1%
Sodium	0.26%	0.0008%
Carbon	41.58%	-
Oxygen	57.67%	-

- = not reported.

Tualang honey: Malaysian multi-floral jungle honey produced by *Apis dorsata*.
Manuka honey: Australian or New Zealand mono-floral honey produced by *Apis mellifera*.

Use

Cosmetic

The safety of the cosmetic ingredients addressed in this assessment is evaluated based on data received from the US Food and Drug Administration (FDA) and the cosmetics industry on the expected use of these ingredients in cosmetics. Use frequencies of individual ingredients in cosmetics are collected from manufacturers and reported by cosmetic product category in the FDA Voluntary Cosmetic Registration Program (VCRP) database. Use concentration data are submitted by the cosmetic industry in response to a survey, conducted by the Personal Care Products Council (Council), of maximum reported use concentrations by product category.

According to 2020 VCRP survey data, Honey is reported to be used in 1059 formulations (671 of which are leave-on formulations), and Honey Extract is reported to be used in 398

formulations (192 of which are leave-on formulations; Table 5).²⁷ All other in-use ingredients are reported to be used in 6 formulations or less. The results of a 2018 concentration of use survey conducted by the Council indicate Honey also has the highest concentration of use; it is used at up to 22% in paste masks and mud packs (which are considered rinse-off formulations).²⁸ The highest concentration of use reported for leave-on products was in formulations containing Honey Extract at up to 7% in body and hand products. Use concentration data were reported for Honey Cocoates in response to the Council survey (it is used at up to 2% in skin cleansing formulations), but no uses were reported in the VCRP; it should be presumed there is at least one use in a skin cleansing formulation, for which the concentration is reported. Conversely, VCRP data are available for Honey Powder, but concentration of use data were not reported. The ingredients not in use according to the VCRP and industry survey are Hydrolyzed Honey and Hydrolyzed Honey Protein.

Honey is reported to be used in baby products, products that would be used near the eye, and products that could result in incidental ingestion and mucous membrane exposure. Honey is reported to be used in 13 baby products and at up to 0.01%. It is also reported to be used in 20 lipstick formulations (up to 3%), 1 dentifrice formulation (up to 0.00035%), 5 “other” oral hygiene product formulations (up to 0.1%), and 1 mouthwash and breath freshener formulation (concentration unknown). Honey could also result in mucous membrane exposure as it is used at up to 3% in bath soaps and detergent formulation.

Additionally, Honey and Honey Extract are used in cosmetic sprays and could possibly be inhaled; for example, Honey is reported to be used in colognes and toilet waters and in hair sprays at up to 0.25% and 0.1%, respectively. In practice, most droplets/particles released from cosmetic sprays have aerodynamic equivalent diameters >10 µm, with propellant sprays yielding a greater fraction of droplets/particles <10 µm compared with pump sprays.^{29,30} Therefore, most droplets/particles incidentally inhaled from cosmetic sprays would be deposited in the nasopharyngeal and thoracic regions of the respiratory tract and would not be respirable

Table 5. Frequency (2020) and Concentration (2018) of Use of Honey Ingredients.^{27,28}

	# of Uses	Max Conc of Use (%)	# of Uses	Max Conc of Use (%)	# of Uses	Max Conc of Use (%)
	Honey		Honey Cocoates		Honey Extract	
Totals ^a	1059	0.00001–22	NR	2	398	0.000002–7
Duration of use						
Leave-on	671	0.0001–3	NR	NR	192	0.0000034–7
Rinse-off	377	0.00001–22	NR	2	191	0.000002–0.01
Diluted for (bath) use	11	NR	NR	NR	15	NR
Exposure type						
Eye area	23	3	NR	NR	10	NR
Incidental ingestion	27	0.00035–3	NR	NR	17	NR
Incidental inhalation-spray	2; 186 ^b ; 353 ^c	0.001–0.25 0.01–0.75 ^c	NR	NR	2; 70 ^b ; 73 ^c	0.0000034–0.001
Incidental inhalation-powder	186 ^b ; 7 ^d	3; 0.0005–3 ^d	NR	NR	70 ^b	0.0001; 0.001–7 ^d
Dermal contact	882	0.00001–22	NR	NR	291	0.000002–7
Deodorant (underarm)	NR	NR	NR	NR	NR	NR
Hair - non-coloring	145	0.000039–10	NR	NR	62	0.0000034–0.005
Hair-coloring	2	0.01–0.8	NR	NR	25	0.0001–0.006
Nail	1	NR	NR	NR	NR	NR
Mucous membrane	209	0.00035–3	NR	NR	89	0.00028–0.01
Baby products	13	0.01	NR	NR	NR	0.00051
	Honey Powder		Hydrogenated Honey			
Totals ^a		6		NR	6	0.25
Duration of use						
Leave-on		3		NR	1	NR
Rinse off		3		NR	5	0.25
Diluted for (bath) use		NR		NR	NR	NR
Exposure type						
Eye area		NR		NR	NR	NR
Incidental ingestion		NR		NR	NR	NR
Incidental inhalation-spray		2 ^b		NR	1 ^b	NR
Incidental inhalation-powder		2 ^b		NR	1 ^b	NR
Dermal contact		6		NR	6	0.25
Deodorant (underarm)		NR		NR	NR	NR
Hair - non-coloring		NR		NR	NR	NR
Hair-coloring		NR		NR	NR	NR
Nail		NR		NR	NR	NR
Mucous membrane		NR		NR	NR	NR
Baby products		NR		NR	NR	NR

^aBecause each ingredient may be used in cosmetics with multiple exposure types, the sum of all exposure types may not equal the sum of total uses.

^bNot specified whether a spray or a powder, but it is possible the use can be as a spray or a powder, therefore the information is captured in both categories.

^cIt is possible these products are sprays, but it is not specified whether the reported uses are sprays.

^dIt is possible these products are powders, but it is not specified whether the reported uses are powders.

Abbreviation: NR: no reported use.

(i.e., they would not enter the lungs) to any appreciable amount.^{30,31} Honey is reportedly used in face powders at concentrations up to 3%, and could possibly be inhaled. Honey Extract is also reported to be used in powders (dusting and talcum) at up to 0.0001%. Conservative estimates of inhalation exposures to respirable particles during the use of

loose powder cosmetic products are 400-fold to 1000-fold less than protective regulatory and guidance limits for inert airborne respirable particles in the air.^{32–34}

The honey-derived ingredients in this report are not restricted from use in any way under the rules governing cosmetic products in the European Union.³⁵

Non-Cosmetic

Food. Raw honey has been consumed worldwide for centuries.³⁶ Honey is commonly used as a sweetener and flavoring agent in many foods. Honey is listed in the US Environmental Protection Agency (EPA) Inert Finder Database as approved for food and non-food use pesticide products.³⁷ For food use, it is regulated under 40 CFR 180.950a. In addition, the FDA requires proper labeling of honey and honey products to ensure that these products are not adulterated and misbranded. All honey and honey products must be labeled in accordance with sections 402 and 403 of the Federal Food, Drug, and Cosmetic Act (21 USC 342 and 343). The international FAO/WHO Codex Alimentarius Standard requires that:³⁸

Honey sold as such shall not have added to it any food ingredient, including food additives, nor shall any other additions be made other than honey. Honey shall not have any objectionable matter, flavor, aroma, or taint absorbed from foreign matter during its processing and storage. The honey shall not have begun to ferment or effervesce. No pollen or constituent particular to honey may be removed except where this is unavoidable in the remove of foreign inorganic or organic matter. Honey shall not be heated or processed to such an extent that its essential composition is changed and/or quality is impaired.

Although rare, infant botulism has been reported after ingestion of honey due to *Clostridium botulinum* spores.³⁹ Because of this, the FDA, the Centers for Disease Control and Prevention, and the American Academy of Pediatrics, recommend not feeding honey to infants younger than 12 months. Neither *Clostridium botulinum* spores nor the neurotoxins are able to penetrate the skin; however, damaged skin may be affected.⁴⁰

Medicine. Honey can be found as an ingredient in over-the-counter (OTC) cough and cold medications.⁴¹ Currently, there is an FDA-approved dermal dressing containing manuka or *Leptospermum* honey, used for the management of wounds and burns.⁴² Examples of wounds that are treated with this dressing are diabetic foot ulcers, leg ulcers, pressure ulcers, partial thickness burns, and surgical wounds.⁴³

Traditionally, honey has been used as an antibacterial, antiseptic, anti-inflammatory, and apitherapeutic agent.³⁶ Honey is commonly used for treatment of cuts, eczema, dermatitis, skin diseases, Fournier's gangrene, burns, ulcers, surgical wounds, fungating wounds, pressure sores, and cancer or broken skin.⁴⁴ Traditional, Ayurvedic treatments utilize honey for cardiac pain, palpitations, and eye ailments.⁴⁵

Toxicokinetic Studies

Toxicokinetic studies were not available regarding these honey derived-ingredients. However, toxicokinetic information on some of the relevant, primary components of honey

(fructose, glucose, and maltose) can be found in the Panel's report on monosaccharides and disaccharides.⁴

Toxicological Studies

No general toxicological studies were found in the published literature, and unpublished data were not submitted.

Developmental and Reproductive Toxicity Studies

The effect of Palestinian honey on spermatogenesis was studied in male albino rats (12 rats/group) after 20 days of treatment.⁴⁶ Group A was given a 5% solution of Palestinian honey in drinking water, group B was treated with 5% sucrose in drinking water, and group C served as the control group and was given untreated drinking water. No significant effects on total body weight or weights of the testis, seminal vesicles, spleen, kidneys, liver, heart, or brain were noted. Rats treated with Palestinian honey displayed a significant increase in epididymal sperm count by 37% ($P \leq 0.05$). The activity of testicular marker enzymes for spermatogenesis such as sorbitol dehydrogenase was increased by 31%, and lactate dehydrogenase was reduced by 48%, indicating an induction of spermatogenesis.

A study was performed in order to examine the effect of honey on the reproductive system of rat male offspring.⁴⁷ Dams were divided into 10 rats/group. The control group received no treatment while treated animals were given honey (0.2 g/kg bw), daily, from day 1 of pregnancy to day 10, via gavage. In male offspring, testosterone levels were significantly lower in the treated group compared to the control group. Sperm counts, follicle stimulating hormone levels, and testes/epididymis weights were similar in control and honey-treated groups. The percentage of abnormal sperm was significantly higher in offspring of dams treated with honey compared to the control group.

Genotoxicity Studies

No genotoxicity studies were found in the published literature, and unpublished data were not submitted.

Anti-Mutagenicity

The potential anti-mutagenic effect of various honeys (fireweed, tupelo, Hawaiian Christmas berry, clover, acacia, buckwheat, and soybean) on 3-amino-1,4-dimethyl-5H-pyrido[4,3-b]indole (Trp-p-1), was studied.⁴⁸ Trp-p-1 is a commonly encountered food mutagen, and has been demonstrated to be mutagenic in bacteria and carcinogenic in animals. The anti-mutagenic effects of the honeys were assayed according to an Ames assay, with slight modification. All assays were performed in a final volume of 1 ml containing potassium phosphate buffer, Trp-p-1 (5 μ l of 20 μ g/ml in dimethyl

sulfoxide), 4% S9 mix (500 μ l), test strain *Salmonella typhimurium* TA98 (2×10^{10} cells/ml), and different honey solutions. Acacia, fireweed, soy, and tupelo honeys demonstrated enhanced anti-mutagenicity above 1 mg/ml, with inhibition between 40.3 and 62.9%; concentrations above 20 mg/ml did not further enhance anti-mutagenic effects. Clover and Hawaiian Christmas berry honey were most effective at 20 mg/ml, with 64.8 and 59.6% inhibition, respectively. The greatest inhibitory effect of buckwheat honey was observed at 1 mg/ml (52.1%).

Carcinogenicity Studies

No carcinogenicity studies were found in the published literature, and unpublished data were not submitted.

Anti-Carcinogenicity Studies

The potential anti-carcinogenic effect of tualang honey on breast cancer was studied in rats.⁴⁹ Forty female Sprague-Dawley rats were given 80 mg/kg 7,12-dimethylbenz[a]anthracene (DMBA) via gavage. Rats were then divided into four groups. Animals in group 1 were given only distilled water. Animals in groups 2, 3, and 4 were given 0.2, 1, and 2 g/kg bw/d tualang honey diluted in 0.5 ml water, respectively, via gavage, for 150 days. After treatment, animals were euthanized. Breast cancers in the honey-treated groups had smaller tumor size compared to controls. In addition, the number of cancers developed in honey-treated rats was significantly lower than control groups ($P < .05$). The majority of the cancers in the control groups were high grade, while cancers in honey-treated groups were of medium- or low-grade. These effects, however, were not dose-dependent.

Anti-Tumorigenicity

The anti-tumoral therapeutic effects of tualang honey and manuka honey was studied in rats (10/group).⁵⁰ Thirty female Sprague-Dawley rats were given an 80 mg/kg injection of the carcinogen 1-methyl-1-nitrosourea (MNU); an additional 10 female rats were left untreated. Treatment with honey started when the first tumor reached 10–12 mm in size. Positive (tumor induction and no honey treatment) and negative controls (no tumor induction or honey treatment) were included. Treatment groups were fed either tualang or manuka honey (1 g/kg bw/d) for 120 days. On the 120th day of treatment, rats were euthanized. Rats in the positive control group had the highest median number of breast tumors compared to groups treated with either honey. Groups treated with honey showed a significant reduction in tumor size and weight compared to the positive control group. The percent reduction in the size of primary tumors was greater with tualang honey (70.82%), as compared to manuka honey (57%). Tumor masses in the positive control group were solid, large in size, and hard in consistency, exhibiting areas of

necrosis and hemorrhage. Both honey-treated groups had tumors which were softer, paler, and smaller in size. Tumors in the positive control group were observed to have increased heterogenous nuclei formation, which were hyperchromatic, vesicular, and highly pleomorphic, with moderate cytoplasm increased mitotic activity compared with the honey-treated groups, which had fatty tissue, small nuclei, and cystic spaces.

Other Relevant Studies

Airway Inflammation Reduction

New Zealand white rabbits (5/group) were dosed twice with an intraperitoneal injection of ovalbumin (OVA) and aluminum hydroxide on days 1 and 14.⁵¹ Tualang honey was then given via a nebulizer from days 23 to 25 at concentrations of either 25 or 50%, diluted in sterile phosphate buffer saline (5 ml for 20 min). After treatment with aerosolized honey, animals were either euthanized, or, further exposed to aerosolized OVA for 3 days starting from day 28 and euthanized on day 31. The effects of honey on the inflammatory cell response, airway inflammation, and goblet cell hyperplasia were assessed. Treatment with aerosolized honey reduced the number of airway inflammatory cells present in bronchioalveolar lavage fluid and inhibited goblet cell hyperplasia. In addition, treatment with aerosolized honey led to a significant decrease in the thickening of the epithelial and mucosal regions.

Nasal Respiratory Mucosa

A study was performed in New Zealand white rabbits (2/group) to evaluate the effect of manuka honey on nasal respiratory mucosa.⁵² The left nasal cavity of each rabbit was irrigated once daily with 1.5 ml of a 33% mixture of manuka honey with saline; groups were treated for either 3, 7, or 14 consecutive days, and then euthanized. The last group was treated for 14 days followed by a 14-day washout period, and then euthanized the following morning. The right nasal cavity of each rabbits served as a control, and was not treated. The mucosa were examined by light microscopy. No histological evidence of inflammation, mucosal injury, or significant morphological changes were observed.

Allergic Potential Following Ingestion

Twenty subjects participated in a 12-week study to determine the allergic potential of manuka and multi-floral honey.⁵³ The participants ate a normal diet with the inclusion of the allocated honey. For the first 2 weeks, all honey was excluded from the diet; then, participants consumed 20 g honey per day in 2 doses of 10 g each. After 4 weeks, there was another 2-week “washout” period, and the groups swapped to the other type of honey for 4 weeks. Fasting blood samples were collected at the beginning of the study, starting with the first

sample after the initial 2-week washout, and then weekly during the 4-week interventions with honey. Immunoglobulin E (IgE) measurements were carried out on frozen serum collected weekly during each of the honey interventions. IgE levels remained at a level consistent with a non-atopic response during the course of the study. The authors concluded that this level of consumption of manuka and multi-floral honey had no significant effect on allergic status.

Cytotoxicity of Honey-Impregnated Wound Dressing

The potential cytotoxic effect of honey-impregnated wound dressings on human skin keratinocytes and dermal fibroblasts was studied.⁵⁴ Five and 21 days after initiating the tissue culture, the honey-impregnated wound dressing was introduced directly onto the cells in the test wells to allow for cell growth. Small blocks of commercial dressings were then inserted into the wells, adjacent and distal to the tissue explants. The amount of test material used was not stated. Keratinocytes and fibroblasts treated with honey implants displayed a modest uniform increase in early cell proliferation and cell counts per mm. Nuclear and cytoplasmic networks appeared normal, and cell proliferation was also evident immediately adjacent to the product. No cell toxicity was observed.

Cytotoxicity in Cancer Cells

Renal cell carcinoma cells (ACHN) were cultured in a medium containing 10% fetal bovine serum and 5, 10, or 15% honey for 3 consecutive days.⁵⁵ Cell viability was determined by the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay, and apoptotic cells were determined using annexin V-fluorescein isothiocyanate (FITC) by flow cytometry. Honey decreased cell viability and induced apoptosis in malignant cells in a concentration- and time-dependent manner ($P < .001$). The half maximal inhibitory concentration (IC_{50}) values at 48 and 72 h were 1.7 ± 0.04 and 2.1 ± 0.03 $\mu\text{g/ml}$, respectively ($<2.5\%$ honey in each case).

A similar study was performed on human breast cancer (MCF-7, MDA-MB-231), immortalized cervical cancer (HeLa), and normal breast epithelial cells.⁵⁶ Cells were plated at a concentration of 1×10^5 cells/well. The cells were allowed to adhere overnight, and the culture medium was replaced with fresh assay medium supplemented with 2% fetal bovine serum. Cells were then treated with different concentrations of tualang honey (1–10%), and incubated for up to 72 h. Tualang honey induced a statistically significant increase in cell death in MDA-MB-231, MCF-7, and HeLa cancer cell lines in a dose- and time-dependent manner. Treatment of the normal breast epithelial cell line did not show a clear cytotoxic effect, even after 72 h of incubation. Flow cytometric analysis of cells stained with annexin V-FITC and propidium iodide showed that tualang honey significantly increased apoptosis in all cancer cell lines compared to untreated cells.

Dermal Irritation and Sensitization Studies

Human

Honey Extract. A human repeated insult patch test (HRIPT) was performed on 112 subjects using a cosmetic product containing 7% Honey Extract.⁵⁷ Approximately 0.2 ml of the test substance was applied to the upper back, under an occlusive patch. Patches were allowed to remain in direct skin contact for a period of 24 h. Applications were made to the same site, three times a week, for a total number of 9 applications during the induction period. After a 2-week rest period, challenge patches were applied to previously untreated test sites. After 24 h, patches were removed and test sites were evaluated. The test substance did not demonstrate a potential for eliciting dermal irritation or sensitization.

According to a summary report, an HRIPT was performed on 116 subjects using a product containing 0.01% Honey Extract according to the same procedure as above.⁵⁸ The product was tested at a 1% dilution in water (effective test concentration, 0.0001% Honey Extract). Seven individuals displayed low-level reactions (mild erythema) during the induction phase, and one individual displayed a high-level reaction in the induction phase. Eight individuals displayed low-level reactions during the challenge phase. (Individual subject scores were not provided). The test substance was considered by the researchers to be non-sensitizing.

Ocular Irritation Studies

Human

Use Study. A prospective, randomized, paired-eye, investigator-masked trial was performed on 25 subjects to determine the clinical safety of manuka honey eye cream on patients with blepharitis.⁵⁹ The cream (approximately 0.034 g \pm 0.001 g) was placed on the periocular surface of the closed upper and lower eyelids of the affected eye. Applications occurred once a day, at night, for 2 weeks. The untreated eye served as a control. A questionnaire was given to grade the severity of dry eye symptomatology at baseline, and a telephone interview was conducted following the first day of cream application to check for immediate tolerability issues or adverse events. Clinical assessments were performed at baseline, day 7, and day 14 of the treatment period. There were no statistically significant differences in baseline clinical or impression cytology measurements between treated and control eyes. Twenty-three of 25 participants did not report any tolerability issues or adverse effects following the first day of product application. In two individuals, application too close to the eyelash margin and the use of excessive cream was presumed to result in a transient stinging sensation. Irrigation with water and reapplication of a modest quantity of cream resolved the issue in both cases. No other adverse effects were reported throughout the study.

Clinical Studies

Effect on Damaged Pediatric Skin

Eight pediatric patients ranging from 8 months to 13 years of age were evaluated in this study.⁶⁰ Five of the children had second-degree burns, and three had necrotic ulcers, circular skin lesions, and deep cervical trauma. Each child was treated with povidone iodine (10% solution), fusidic acid, and systemic antibiotics, followed by a honey-based ointment. After this initial treatment, patients were instructed to apply honey-containing ointment as well as a dressing impregnated with a polymer containing 20% medical grade honey, daily. The duration and amount of product used in this study were not stated. No adverse effects or allergic reactions were observed.

Case Studies

Anaphylaxis. A 40-year-old woman was referred to a clinic after suspected allergy to honey.⁶¹ At the age of 36, she had two episodes of generalized urticaria 20 min after ingestion of foods with honey. At the age of 37, 5 min after an inadvertent contact with a teaspoon with traces of honey, the patient reported swollen lips, urticaria, and angioedema. After treatment with oral corticosteroids and antihistamines, symptoms were resolved. Skin prick tests with standard panel of extracts from aeroallergens and common allergenic foods yielded negative results. Prick-to-prick tests (PPT) were performed with the previously consumed honey, and eight other kinds of honey (eucalyptus, sunflower, orange-tree, Arbutus-tree, French lavender, heather, flower incense, and rosemary). Results were positive for all honey types. Thirty minutes after the administration of the PPT, the patients suffered from anaphylaxis, generalized urticaria, swollen lips, tongue, and uvula, and hypotension. The same PPT was performed with these honeys in 6 control volunteers (3 healthy individuals, and 3 atopic with pollen sensitization and rhinitis). None of the volunteers displayed a positive skin reaction.

Epicutaneous Sensitization. A 48-year-old woman had been washing her body and hair with products blended with edible honey, and she applied honey to the face as a face pack.⁶² After 8 years of use, the woman developed itching and redness on facial skin as well as conjunctival hyperemia following the use of the face pack containing honey. After washing her body with honey-containing soap, the subject reported urticarial symptoms on her extremities and un-exposed face. One year later, the subject developed abdominal pain and distention after eating yogurt with honey. The patient had positive results for honey-antigen specific IgE antibodies in serum uric acid (UA), equivalent to 1.44 UA/ml, but not for honey bee venoms or Api m 10 (*Apis mellifera* venom component). Results for specific IgE against three cross-reactive carbohydrate determinant marker allergens were negative. Prick tests with honey

gave positive results. Fifteen minutes after oral challenge with 30 ml of honey, the patient developed eyelid swelling, abdominal pain, and oral tingling.

Summary

The 7 honey-derived ingredients in this report all are reported to function in cosmetics as skin-conditioning agents. Other reported cosmetic functions include flavoring agent, abrasive, binder, depilating agent, exfoliant, hair-conditioning agent, and nail-conditioning agent. Honey derived for cosmetic purposes is reported to be produced by the honeybee species *Apis mellifera*, *Tetragonisca angustula*, *Scaptotrigona pectoralis*, and *Melipona becheii*.

Of the ingredients included in this report, Honey has the most reported uses, with a total of 1059; 671 of these are leave-on products. Honey Extract has the second greatest number of overall uses, with a total of 398 (192 are in leave-on formulations). Honey has the highest concentration of use, and is used at up to 22% in paste and mud packs. The highest concentration of use reported for leave-on products was in body and hand products containing Honey Extract at up to 7%. The ingredients not in use according to VCRP data and the industry survey are Hydrolyzed Honey and Hydrolyzed Honey Protein.

Honey is common in food and food products worldwide. Honey can be found in OTC cough and cold medications. Traditional medicine suggests the use of honey for various ailments and skin issues. Currently, there is an FDA-approved dermal dressing containing honey used for the management of wounds and burns.

The effect of Palestinian honey on spermatogenesis was studied in male albino rats. Rats treated with Palestinian honey displayed a significant increase in epididymal sperm count. The activity of testicular marker enzymes for spermatogenesis, such as sorbitol dehydrogenase, was increased, and lactate dehydrogenase was reduced, indicating an induction of spermatogenesis. The effect of honey on the reproductive system of rat male offspring was studied. Testosterone levels were significantly lower in the male offspring of treated animals, compared to control animals. The percentage of abnormal sperm was significantly higher in the offspring of dams treated with honey versus the control group. All other parameters were similar between treated and control group.

The potential anti-mutagenic effect of various honeys on Trp-p-1 was studied. Acacia, fireweed, soy, and tupelo honeys demonstrated enhanced antimutagenicity above 1 mg/ml, with inhibition between 40.3 and 62.9%. Concentrations above 20 mg/ml demonstrated no enhancement of the antimutagenic effects. Clover and Hawaiian Christmas berry honey were most effective at 20 mg/ml, with 64.8 and 59.6% inhibition, respectively. The greatest inhibitory effect of buckwheat honey was observed at 1 mg/ml (52.1%).

In an anti-tumorigenicity study, Sprague-Dawley rats were given an injection of the carcinogen MNU and either given no treatment or treatment with manuka or tualang honey (1 g/kg bw/d) via diet. Groups treated with honey showed a significant reduction in tumor size and weight compared to the nontreated positive control. In addition, breast tumors in the positive control were large and hard, while tumors in honey-treated groups were small and soft.

In New Zealand white rabbits pre-treated with ovalbumin, treatment with aerosolized honey reduced the number of airway inflammatory cells present in bronchioalveolar lavage fluid and inhibited goblet cell hyperplasia. In addition, treatment with aerosolized honey led to a significant decrease in the thickening of the epithelial and mucosal regions. The nasal cavities of New Zealand white rabbits were irrigated with a honey and saline solution. No histological evidence of inflammation, epithelial injury, or significant morphological changes were observed.

Twenty subjects were used in a study to determine the allergic potential of manuka and multi-floral honey following ingestion. IgE levels remained at a level consistent with a non-atopic response during the course of the study.

The potential cytotoxic effect of honey-impregnated wound dressings on human skin keratinocytes and dermal fibroblasts was studied. Keratinocytes and fibroblasts treated with honey implants displayed a modest uniform increase in early cell proliferation and cell counts per mm. No cytotoxic effects were observed.

The anti-carcinogenic potential of honey (up to 15%) was studied using renal cell carcinoma cell lines. Honey decreased cell viability and induced apoptosis in malignant cells in a concentration- and time-dependent manner. A similar study was performed using tualang honey (1–10%) on human breast cancer, cervical cancer, and normal breast epithelial cell lines. Treatment with honey induced cell death in all cancer cell lines, but no clear cytotoxic effect was observed in the normal breast epithelial cells. In a different study, the effect of tualang honey (0.2–2 g/kg) on breast cancer-induced rats was observed. Smaller tumors were observed in honey-treated rats compared to control animals. In addition, the number of cancers developed in honey-treated rats was significantly lower than control groups.

An HRIPT that was performed on 112 subjects using a cosmetic product containing 7% Honey Extract applied using occlusive conditions yielded negative results. In an HRIPT performed on 116 subjects using a test substance containing 0.01% Honey Extract, tested as a 1% dilution, the test substance was considered to be non-sensitizing.

Twenty-five subjects were used in a 2-week prospective study to determine the safety of manuka honey eye cream on blepharitis patients. Twenty-three of 25 participants did not report any tolerability issues or adverse effects following the first day of product application. In two individuals, application too close to the eyelash margin and the use of excessive cream was presumed to result in a transient stinging sensation.

Honey-based ointment was used as part of a treatment of damaged skin in 8 children. No adverse effects or allergic reactions were reported after treatment.

Generalized urticaria was reported in a patient 20 min after ingesting foods with honey. After inadvertent contact with traces of honey, the same patient reported other allergic symptoms. Skin prick tests on common allergenic foods yielded negative results, however, prick-to-prick testing using different honey types yielded positive results.

After using cosmetics blended with edible honey and using honey as a face pack for 8 years, a woman reported allergic reactions after using a face pack containing honey and body soap containing honey. A year after these symptoms occurred, the patient experienced abdominal pain after ingestion of honey. Prick tests with honey yielded positive results.

Discussion

The Panel reviewed the available, relevant data to assess the safety of these honey-derived ingredients as used in cosmetics in the present practices of use and concentration. The Panel noted the lack of sensitization data for six of the seven ingredients, but determined that the available sensitization data on Honey Extract could be used to support the safety of the remaining ingredients. The safety of these ingredients is further supported by historical food use and use in wound dressings.

As Honey Powder in foods has been reported to be 50% filler material, the Panel also discussed the safety of those possible filler ingredients (starch, carboxymethyl cellulose, gum Arabic, maltodextrin, and gelatin), and determined that these ingredients are not of concern. All of the named fillers have been previously reviewed by the Panel, and were considered safe as used in cosmetics.

The Panel expressed concern regarding pesticide residues and endotoxins that may be present in these ingredients. They stressed that the cosmetics industry should continue to use current good manufacturing practices (cGMPs) to limit these impurities. In addition, the Panel noted the importance of avoiding the use of honey derived from toxic plant sources for use in cosmetic formulations.

The Panel discussed the issue of incidental inhalation exposure from formulations that may be aerosolized (e.g., colognes and toilet waters at up to 0.25% Honey). The Panel noted that in aerosol products, most droplets/particles would not be respirable to any appreciable amount. Furthermore, droplets/particles deposited in the nasopharyngeal or bronchial regions of the respiratory tract present no toxicological concerns based on the properties of these ingredients. Coupled with the small actual exposure in the breathing zone and the concentrations at which the ingredients are used, the available information indicates that incidental inhalation would not be a significant route of exposure that might lead to local respiratory or systemic effects. Respiratory safety of these

ingredients was further supported by the lack of negative effects observed in a nasal irrigation and an inhalation study performed in rabbits. A detailed discussion and summary of the Panel's approach to evaluating incidental inhalation exposures to ingredients in cosmetic products is available at <https://www.cir-safety.org/cir-findings>.

Conclusion

The Expert Panel for Cosmetic Ingredient Safety concluded that the following honey-derived ingredients are safe in cosmetics in the present practices of use and concentrations described in the safety assessment.

Honey

Honey Cocoates

Honey Extract

Honey Powder

Hydrogenated Honey

Hydrolyzed Honey*

Hydrolyzed Honey Protein*

*Not reported to be in current use. Were ingredients in this group not in current use to be used in the future, the expectation is that they would be used in product categories and at concentrations comparable to others in this group.

Author's Note

Unpublished sources cited in this report are available from the Director, Cosmetic Ingredient Review, 555 13th Street, NW, Suite 300W, Washington, DC 20004, USA.

Author Contributions

The articles in this supplement were sponsored by the Cosmetic Ingredient Review.

Declaration of Conflicting Interest

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The articles in this supplement were sponsored by the Cosmetic Ingredient Review. The Cosmetic Ingredient Review is financially supported by the Personal Care Products Council.

References

1. Nikitakis J, Kowcz A. *wINCI: International Cosmetic Ingredient Dictionary and Handbook*. Washington, DC: Personal Care Products Council. <https://webdictionary.personalcarecouncil.org/jsp/Home.jsp>. Last Updated: 2019. Accessed June 25, 2019.
2. Burnett C, Fiume M, Bergfeld W, et al. Safety assessment of plant-derived fatty acid oils. *Int J Toxicol*. 2017;36:51S-129S.
3. Alvarez-Suarez J, Giampieri F, Battino M. Honey as a source of dietary antioxidants: structures, bioavailability, and evidence of protective effects against human chronic diseases. *Curr Med Chem*. 2013;20(5):621-638.
4. Fiume M, Bergfeld W, Belsito D, et al. Safety assessment of monosaccharides, disaccharides, and related ingredients as used in cosmetics. *Int J Toxicol*. 2019;38:5S-38S.
5. Chemical and Functional Properties of Food Components Series. In Tomasik P, ed. *Chemical and Functional Properties of Food Saccharides*. Boca Raton, FL: CRC Press, Taylor and Francis Group; 2004:73-79.
6. Satarupa R, Subha G. Physical, chemical, and antioxidant properties of honey: a review. *Asian J Chem Pharm Res*. 2014;2(1):96-99.
7. Anonymous. Honey extract: manufacturing, heavy metals, impurities, allergen, and chemical and physical property information. (Unpublished data submitted by Personal Care Products Council on October 15, 2019); 2019.
8. National Honey Board. Honey floral source guide. Longmont, CO. https://www.bjcp.org/mead/floral_guide.pdf. Last Updated: 2019. Accessed August 14, 2019.
9. Cridland J, Tsutsui N, Ramírez SR. The complex demographic history and evolutionary origin of the Western honey bee, *Apis mellifera*. *Genome Biol Evol*. 2017;9(2):457-472.
10. Batista M, Ramalho M, Soares A. Nesting sites and abundance of Meliponini (Hymenoptera: Apidae) in heterogeneous habitats of the Atlantic Rain Forest, Bahia, Brazil. *Lundiana*. 2003;4(1):19-23.
11. León A, Arias-Castro C, Rodríguez-Mendiola M, Meza-Gordillo R, Gutiérrez-Miceli F, Nieh J. Colony foraging allocation is finely tuned to food distance and sweetness even close to a bee colony. *Entomol Exp Appl*. 2015;155:47-53.
12. Quezada-Euán J, Paxton R, Palmer K, Itzá W, Tay W, Oldroyd B. Morphological and molecular characters reveal differentiation in a Neotropical social bee, *Melipona beecheii* (Apidae: Meliponini). *Adipologie*. 2007;38:247-258.
13. Eyer M, Neumann P, Dietemann V. A look into the cell: honey storage in honey bees, *Apis mellifera*. *PLoS One*. 2016;11(8):e0161059.
14. Ediriweera E, Premarathna N. Medicinal and cosmetic uses of Bee's Honey - a review. *Ayu*. 2012;33(2):178-182.
15. Solabia Group. Manufacturing process: Honey Milk 1.5PS (trade name mixture containing 10.6% Honey Extract). (Unpublished data submitted by the Personal Care Products Council on September 10, 2019); 2012.
16. Solabia Group. Manufacturing process: Glycolysat® of Honey UP (trade name mixture containing 16.5% Honey Extract). (Unpublished data submitted by Personal Care Products Council on September 10, 2019); 2009.
17. Nurhadi B, Andoyo R, Mahani IR. Study the properties of honey powder produced from spray drying and vacuum drying method. *Int Food Res J*. 2012;19(3):907-912.
18. Chua LS, Lee J, Chan G. Honey protein extraction and determination by mass spectrometry. *Anal Bioanal Chem*. 2013;405(10):3063-3074.

19. Istasse T, Jacquet N, Berchem T, Haubruge E, Nguyen B, Richel A. Extraction of honey polyphenols: method development and evidence of *Cis* isomerization. *Anal Chem Insights*. 2016;11: 49-57.
20. Islam MN, Khalil I, Islam A, Gan S. Toxic compounds in honey. *J Appl Toxicol*. 2014;34(7):733-742.
21. Bauer L, Kohlich A, Hirschwehr R, et al. Food allergy to honey: pollen or bee products? *J Allergy Clin Immunol*. 1996;97(1): 65-73.
22. Ahmed A, Othman N. Review of the medicinal effects of tualang honey and a comparison with Manuka honey. *Malays J Med Sci*. 2013;20(3):6-13.
23. Alleva R, Manzella N, Gaetani S, et al. Organic honey supplementation reverses pesticide-induced genotoxicity by modulating DNA damage response. *Mol Nutr Food Res*. 2016; 60(10):2243-2255.
24. Al-Waili N, Salom K, Al-Ghamdi A, Ansari M. Antibiotic, pesticide, and microbial contaminants of honey: human health hazards. *Sci World J*. 2012;2012:930849. doi:10.1100/2012/930849. Last Updated: 2012. Accessed August 15, 2019.
25. Joint FAO/WHO Food Standard Programme Codex Alimentarius Commission. Report of the Seventh Session of the Codex Committee on Sugars. London, UK; 2000. https://www.fao.org/input/download/report/277/A101_25c.pdf. Accessed August 8, 2019.
26. The Council of the European Union. Council directive 2001 of 20 December 2001/110/EC relating to honey; 2002:47-52.
27. U.S. Food and Drug Administration Center for Food Safety & Applied Nutrition (CFSAN). *Voluntary Cosmetic Registration Program - Frequency of Use of Cosmetic Ingredients*. College Park, MD; 2020. ((Obtained under the Freedom of Information Act from CFSAN; requested as "Frequency of Use Data" January 6, 2020; received January 13, 2020)).
28. Personal Care Products Council. Concentration of use by FDA product category: honey ingredients. (Unpublished data submitted to Personal Care Products Council on October 12, 2018); 2018.
29. Johnsen M. The influence of particle size. *Spray Technol Market*. 2004;14(11):24-27.
30. Rothe H, Fautz R, Gerber E, et al. Special aspects of cosmetic spray safety evaluations: principles on inhalation risk assessment. *Toxicol Lett*. 2011;205(2):97-104.
31. Bremmer HJ, Prud'homme de Lodder L, van Engelen J. Cosmetics Fact Sheet: to assess the risks for the consumer; 2006, 1-77. <https://www.rivm.nl/bibliotheek/rapporten/320104001.pdf>. Updated version for ConsExpo4. Accessed June 25, 2019.
32. CIR Science and Support Committee of the Personal Care Products Council (CIR SCC). (Nov 3rd) cosmetic powder exposure. Unpublished data submitted by the Personal Care Products Council; 2015.
33. Aylott R, Byrne G, Middleton J, Roberts M. Normal use levels of respirable cosmetic talc: preliminary study. *Int J Cosmet Sci*. 1979;1(3):177-186.
34. Russell R, Merz R, Sherman W, Sivertson JN. The determination of respirable particles in talcum powder. *Food Cosmet Toxicol*. 1979;17(2):117-122.
35. European Commission. CosIng database; following Cosmetic Regulation No. 1223/2009. <https://ec.europa.eu/growth/tools-databases/cosing/>. Last Updated: 2016. Accessed December 07, 2019.
36. Ajibola A. Novel insights into the health importance of natural honey. *Malays J Med Sci*. 2015;22(5):7-22.
37. US Environmental Protection Agency (EPA). https://iaspub.epa.gov/apex/pesticides/f?p=INERTFINDER:3::NO::P3_ID:7081. Last Updated: 2019. Accessed July 15, 2019.
38. Food and Agriculture Organization of the United Nations (FAO) and World Health Organization (WHO). *Standard for Honey CXS 12-1981*. Standards CAIF; 2019.
39. US Food and Drug Administration (FDA). Honey pacifiers suspected in Texas infant botulism cases. <https://www.fda.gov/food/alerts-advisories-safety-information/honey-pacifiers-suspected-texas-infant-botulism-cases>. Last Updated: 2018. Accessed July 18, 2019.
40. Nantel A, World Health Organization. Clostridium botulinum, international programme on chemical safety poisons information monograph 858 bacteria. <https://www.who.int/csr/deliblepidemics/clostridiumbotulism.pdf>. Last Updated: 2002. Accessed August 15, 2019.
41. Goldman R. Honey for treatment of cough in children. *Can Fam Physician*. 2014;60(12):1107-1110.
42. US Food and Drug Administration (FDA). Department of Health and Human Services. 510(k) summary for derma sciences medihoney dressings with active Manuka honey. https://www.accessdata.fda.gov/cdrh_docs/pdf8/K080315.pdf. Last Updated: 2008. Accessed August 15, 2019.
43. Derma Sciences. *Wound & Burn Dressing MEDIHONEY® with Active Leptospermum Honey*. Plainsboro, NJ: Derma Sciences Headquarters. <https://www.dermasciences.com/medihoney>. Last Updated: 2019. Accessed July 07, 2019.
44. McLoone P, Oluwadun A, Warnock M, Fyfe L. Honey: a therapeutic agent for disorders of the skin. *Cent Asian J Glob Health*. 2016;5(1):241.
45. Eteraf-Oskouei T, Najafi M. Traditional and modern uses of natural honey in human diseases: a review. *Iran J Basic Med Sci*. 2013;16(6):731-742.
46. Abdul-Ghani A, Dabdoub N, Muhammad R, Abdul-Ghani R, Qazzaz M. Effect of Palestinian honey on spermatogenesis in rats. *J Med Food*. 2008;11(4):799-802.
47. Haron M, Mohamed M. Effect of honey on the reproductive system of male rat offspring exposed to prenatal restraint stress. *Andrologia*. 2016;48(5):525-531.
48. Wang X, Andrae L, Engeseth N. Antimutagenic effect of various honeys and sugars against Trp-p-1. *J Agric Food Chem*. 2002; 50(23):6923-6928.
49. Kadir E, Sulaiman S, Yahya N, Othman N. Inhibitory effects of Tualang Honey on experimental breast cancer in rats: a preliminary study. *Asian Pac J Cancer Prev*. 2013;14(4): 2249-2254.
50. Ahmed S, Othman N. The anti-cancer effects of Tualang honey in modulating breast carcinogenesis: an experimental animal study. *BMC Complement Altern Med*. 2017;17(1):208.

51. Kamaruzaman N, Sulaiman S, Kaur G, Yahaya B. Inhalation of honey reduces airway inflammation and histopathological changes in a rabbit model of ovalbumin-induced chronic asthma. *BMC Complement Altern Med*. 2014;14:176.
52. Kilty S, Al-Mutari D, Duval M, Groleau M, De Nanassy J, Gomes M. Manuka honey: histological effect on respiratory mucosa. *Am J Rhinol Allergy*. 2010;24(2):247.
53. Wallace A, Eady S, Miles M, et al. Demonstrating the safety of manuka honey UMF 20+ in a human clinical trial with healthy individuals. *Br J Nutr*. 2010;103(7):1023-1028.
54. Du Toit D, Page B. An in vitro evaluation of the cell toxicity of honey and silver dressings. *J Wound Care*. 2009;18(9):383-389.
55. Samarghandian S, Afshari J, Davoodi S. Honey induces apoptosis in renal cell carcinoma. *Pharmacogn Mag*. 2011;7:46-52.
56. Fauzi A, Norazmi M, Yaacob N. Tualang honey induces apoptosis and disrupts the mitochondrial membrane potential of human breast and cervical cancer cell lines. *Food Chem Toxicol*. 2011;49(4):871-878.
57. Chemical Research Laboratories Inc. Repeated insult patch test (RIPT) - Shelanski method (product containing 7% Honey Extract). (Unpublished data submitted by Personal Care Products Council on September 24, 2019); 2015.
58. Anonymous. Summary of an HRIPT (product contains 0.01% honey extract). (Unpublished data submitted by Personal Care Products Council on November 19, 2019); 2019.
59. Craig J, Wang M, Ganesalingam K, et al. Randomised masked trial of the clinical safety and tolerability of MGO Manuka Honey eye cream for the management of blepharitis. *BMJ Open Ophthalmol*. 2017;1(1):e000066.
60. Smaropoulos E, Dimitriadou E, Romeos S. Honey-based therapy for paediatric burns and dermal trauma compared to standard hospital protocol. *Wounds U K*. 2011;7(1):33-40.
61. Aguiar R, Duarte F, Mendes A, Bartolomé B, Barbosa M. Anaphylaxis caused by honey: a case report. *Asia Pac Allergy*. 2017;7(1):48-50.
62. Katayama M, Inomata N, Inagawa N, Fukuro S, Aihara M. A case of contact urticaria syndrome stage 3 after honey ingestion induced by epicutaneous sensitization during skin care with honey. *Contact Dermat*. 2016;74:189-191.